

## 7.3) One-tailed tests

## Worked example

Joan believes a six-sided dice is biased in favour of rolling a 4.

She rolls the dice 10 times and rolls a 4 five times.

Using a 5% significance level, test her belief.

## Your turn

John believes a coin is biased in favour of landing with tails uppermost.

He tosses the coin 8 times and it lands on tails 7 times. Using a 5% significance level, test his belief.

$X$  = number of times coin lands with tails uppermost

$p$  = probability/proportion of times coin lands with tails uppermost

$$H_0: p = 0.5$$

$$H_1: p > 0.5$$

Assume  $H_0$  true.  $X \sim B(8, 0.5)$

5% significance level

Reject  $H_0$  if  $P(X \geq 7) < 0.05$

Test  $P(X \geq 7)$

$$= 1 - P(X \leq 6)$$

$$= 0.0351 \dots < 0.05$$

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to support John's belief that the coin is biased in favour of landing with tails uppermost.

## Worked example

Joan believes a six-sided dice is biased in favour of rolling a 4.

She rolls the dice 10 times and rolls a 4 five times.

- Using a 5% significance level, find the critical region to test her belief.
- Joan rolled a 4 three times. Comment on this observation in light of the critical region.

## Your turn

John believes a coin is biased in favour of landing with tails uppermost.

He tosses the coin 8 times.

- Using a 5% significance level, find the critical region to test his belief.
- The coin landed on tails 7 times. Comment on this observation in light of the critical region.

a)  $X$  = number of times coin lands with tails uppermost  
 $p$  = probability/proportion of times coin lands with tails uppermost

$$H_0: p = 0.5$$

$$H_1: p > 0.5$$

Assume  $H_0$  true.  $X \sim B(8, 0.5)$

5% significance level

Reject  $H_0$  if  $P(X \geq x) < 0.05$

$$1 - P(X \leq x - 1) < 0.05$$

$$-P(X \leq x - 1) < -0.95$$

$$P(X \leq x - 1) > 0.95$$

$$P(X \leq 5) = 0.8554 \dots < 0.95$$

$$P(X \leq 6) = 0.9648 \dots > 0.95$$

$$x - 1 = 6$$

$$\therefore x = 7$$

Critical region: Reject  $H_0$  if  $7 \leq X \leq 8$

b) 7 is in the critical region.

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to support John's belief that the coin is biased in favour of landing with tails uppermost

## Worked example

An election candidate believes he has the support of 30% of the residents in a particular town.

A researcher wants to test, at the 10% significance level, whether the candidate is over-estimating his support.

The researcher asks 30 people whether they support the candidate or not. 6 people say they support the candidate.

Carry out a hypothesis test for the researcher.

## Your turn

An election candidate believes she has the support of 40% of the residents in a particular town.

A researcher wants to test, at the 5% significance level, whether the candidate is over-estimating her support.

The researcher asks 20 people whether they support the candidate or not. 4 people say they support the candidate.

Carry out a hypothesis test for the researcher.

$X$  = number of people who say they support the candidate

$p$  = probability/proportion of people who say they support the candidate

$$H_0: p = 0.4$$

$$H_1: p < 0.4$$

Assume  $H_0$  true.  $X \sim B(20, 0.4)$

5% significance level

Reject  $H_0$  if  $P(X \leq 4) < 0.05$

Test  $P(X \leq 4)$

$$= 0.0509 \dots > 0.05$$

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to suggest the candidate is over-estimating her support.

## Worked example

An election candidate believes he has the support of 30% of the residents in a particular town.

A researcher wants to test, at the 10% significance level, whether the candidate is over-estimating his support.

The researcher asks 30 people whether they support the candidate or not.

- Find the critical region for this test.
- 6 people say they support the candidate. Comment on this observation in light of the critical region.

## Your turn

An election candidate believes she has the support of 40% of the residents in a particular town.

A researcher wants to test, at the 5% significance level, whether the candidate is over-estimating her support.

The researcher asks 20 people whether they support the candidate or not.

- Find the critical region for this test.
- 4 people say they support the candidate. Comment on this observation in light of the critical region.

a)  $X$  = number of people who say they support the candidate

$p$  = probability/proportion of people who say they support the candidate

$$H_0: p = 0.4$$

$$H_1: p < 0.4$$

Assume  $H_0$  true.  $X \sim B(20, 0.4)$

5% significance level

Reject  $H_0$  if  $P(X \leq x) < 0.05$

$$P(X \leq 4) = 0.0509 \dots > 0.05$$

$$P(X \leq 3) = 0.0159 \dots < 0.05$$

$$\therefore x = 4$$

Critical region: Reject  $H_0$  if  $0 \leq X \leq 3$

b) 4 is not in the critical region.

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to suggest the candidate is over-estimating her support.

## Worked example

An election candidate believes he has the support of 30% of the residents in a particular town.

A researcher wants to test, at the 1% significance level, whether the candidate is under-estimating his support.

The researcher asks 30 people whether they support the candidate or not. 14 people say they support the candidate.

Carry out a hypothesis test for the researcher.

## Your turn

An election candidate believes she has the support of 40% of the residents in a particular town.

A researcher wants to test, at the 2% significance level, whether the candidate is under-estimating her support.

The researcher asks 20 people whether they support the candidate or not. 14 people say they support the candidate.

Carry out a hypothesis test for the researcher.

$X$  = number of people who say they support the candidate

$p$  = probability/proportion of people who say they support the candidate

$$H_0: p = 0.4$$

$$H_1: p > 0.4$$

Assume  $H_0$  true.  $X \sim B(20, 0.4)$

5% significance level

Reject  $H_0$  if  $P(X \geq 14) < 0.02$

Test  $P(X \geq 14)$

$$= 1 - P(X \leq 13)$$

$$= 0.00646 \dots < 0.02$$

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to suggest the candidate is under-estimating her support.

## Worked example

An election candidate believes he has the support of 30% of the residents in a particular town.

A researcher wants to test, at the 1% significance level, whether the candidate is under-estimating his support.

The researcher asks 30 people whether they support the candidate or not.

- Find the critical region for this test.
- 14 people say they support the candidate. Comment on this observation in light of the critical region.

## Your turn

An election candidate believes she has the support of 40% of the residents in a particular town.

A researcher wants to test, at the 2% significance level, whether the candidate is under-estimating her support.

The researcher asks 20 people whether they support the candidate or not.

- Find the critical region for this test.
- 14 people say they support the candidate. Comment on this observation in light of the critical region.

a)  $X$  = number of people who say they support the candidate  
 $p$  = probability/proportion of people who say they support the candidate

$$H_0: p = 0.4$$

$$H_1: p > 0.4$$

Assume  $H_0$  true.  $X \sim B(20, 0.4)$

5% significance level

Reject  $H_0$  if  $P(X \geq x) < 0.02$

$$1 - P(X \leq x - 1) < 0.02$$

$$-P(X \leq x - 1) < -0.98$$

$$P(X \leq x - 1) > 0.98$$

$$P(X \leq 12) = 0.9789 \dots < 0.98$$

$$P(X \leq 13) = 0.9935 \dots > 0.98$$

$$x - 1 = 13$$

$$\therefore x = 14$$

Critical region: Reject  $H_0$  if  $14 \leq X \leq 20$

b) 14 is in the critical region.

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to suggest the candidate is under-estimating her support.

## Worked example

In a manufacturing process, the proportion of faulty lightbulbs is, based on historical data, 0.08. The manufacturing process is changed. A sample of 200 lightbulbs is tested. 8 lightbulbs are found to be faulty. The manager wishes to test at the 2% significance level whether or not there has been a reduction in the proportion of faulty lightbulbs. Carry out this hypothesis test.

## Your turn

In a manufacturing process, the proportion of faulty bolts is, based on historical data, 0.07. The manufacturing process is changed. A sample of 100 bolts is tested. 2 bolts are found to be faulty. The manager wishes to test at the 1% significance level whether or not there has been a reduction in the proportion of faulty bolts. Carry out this hypothesis test.

$X$  = number of fault bolts

$p$  = probability/proportion of faulty bolts

$H_0: p = 0.07$

$H_1: p < 0.07$

Assume  $H_0$  true.  $X \sim B(100, 0.07)$

1% significance level

Reject  $H_0$  if  $P(X \leq 2) < 0.01$

Test  $P(X \leq 2)$

$= 0.0257 \dots > 0.01$

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to suggest there has been a reduction in the proportion of faulty bolts.



## Worked example

In a manufacturing process, the proportion of faulty lightbulbs is, based on historical data, 0.08. The manufacturing process is changed. The manager wishes to test at the 2% significance level whether or not there has been a reduction in the proportion of faulty lightbulbs. A sample of 200 lightbulbs is tested.

- Find the critical region for this test.
- 8 lightbulbs are found to be faulty. Comment on this observation in light of the critical region.

## Your turn

In a manufacturing process, the proportion of faulty bolts is, based on historical data, 0.07. The manufacturing process is changed. The manager wishes to test at the 1% significance level whether or not there has been a reduction in the proportion of faulty bolts. A sample of 100 bolts is tested.

- Find the critical region for this test.
- 2 bolts are found to be faulty. Comment on this observation in light of the critical region.

a)  $X =$  number of fault bolts  
 $p =$  probability/proportion of faulty bolts

$$H_0: p = 0.07$$

$$H_1: p < 0.07$$

Assume  $H_0$  true.  $X \sim B(100, 0.07)$

1% significance level

Reject  $H_0$  if  $P(X \leq x) < 0.01$

$$P(X \leq 2) = 0.0257 \dots > 0.01$$

$$P(X \leq 1) = 0.0060 \dots < 0.01$$

$$\therefore x = 1$$

Critical region: Reject  $H_0$  if  $0 \leq X \leq 1$

b) 2 is not in the critical region.

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to suggest there has been a reduction in the proportion of faulty bolts.

## Worked example

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.8% of the time.

They test the benefits of the drug on 4500 patients.

The test is successful in 4498 cases.

Is there enough evidence, at the 1% significance level, to support the medical team's claim?

## Your turn

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.5% of the time.

They test the benefits of the drug on 2500 patients.

The test is successful in 2494 cases.

Is there enough evidence, at the 5% significance level, to support the medical team's claim?

$X$  = number of successful tests

$p$  = probability/proportion of successful tests

$H_0: p = 0.995$

$H_1: p > 0.995$

Assume  $H_0$  true.  $X \sim B(2500, 0.995)$

5% significance level

Reject  $H_0$  if  $P(X \geq 2494) < 0.05$

Test  $P(X \geq 2494) = 0.0342 \dots < 0.05$

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to support the medical team's claim

## Worked example

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.8% of the time.

They test the benefits of the drug on 4500 patients.

The test is successful in 4497 cases.

Is there enough evidence, at the 1% significance level, to support the medical team's claim?

## Your turn

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.5% of the time.

They test the benefits of the drug on 2500 patients.

The test is successful in 2493 cases.

Is there enough evidence, at the 5% significance level, to support the medical team's claim?

$X$  = number of successful tests

$p$  = probability/proportion of successful tests

$H_0: p = 0.995$

$H_1: p > 0.995$

Assume  $H_0$  true.  $X \sim B(2500, 0.995)$

5% significance level

Reject  $H_0$  if  $P(X \geq 2493) < 0.05$

Test  $P(X \geq 2493) = 0.06934 \dots > 0.05$

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to support the medical team's claim

## Worked example

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.8% of the time.

They test the benefits of the drug on 4500 patients.

- Find the critical region for this test at the 1% significance level.
- The test is successful in 4498 cases. Comment on this observation in light of the critical region.

## Your turn

A medical team are testing the effectiveness of a new drug.

They claim that the test is successful more than 99.5% of the time.

They test the benefits of the drug on 2500 patients.

- Find the critical region for this test at the 5% significance level.
- The test is successful in 2493 cases. Comment on this observation in light of the critical region.

a)  $X =$  number of successful tests

$p =$  probability/proportion of successful tests

$$H_0: p = 0.995$$

$$H_1: p > 0.995$$

Under  $H_0$ ,  $X \sim B(2500, 0.995)$

5% significance level

Reject  $H_0$  if  $P(X \geq x) < 0.05$

$$1 - P(X \leq x - 1) < 0.05$$

$$-P(X \leq x - 1) < -0.95$$

$$P(X \leq x - 1) > 0.95$$

$$P(X \leq 2492) = 0.9306 \dots < 0.95$$

$$P(X \leq 2493) = 0.9657 \dots > 0.95$$

$$x - 1 = 2493$$

$$\therefore x = 2494$$

Critical region: Reject  $H_0$  if  $2494 \leq x \leq 2500$

b) 2494 is in the critical region.

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to support the medical team's claim

## Worked example

A medical team are testing the negative side effects of a new drug.  
They claim that the drug gives negative side effects less than 0.2% of the time.  
They test the drug on 4500 patients.  
The drug has negative side effects in 2 patients.  
Is there enough evidence, at the 1% significance level, to support the medical team's claim?

## Your turn

A medical team are testing the negative side effects of a new drug.  
They claim that the drug gives negative side effects less than 0.5% of the time.  
They test the drug on 2500 patients.  
The drug has negative side effects in 3 patients.  
Is there enough evidence, at the 5% significance level, to support the medical team's claim?

$X$  = number of tests with negative side effects  
 $p$  = probability/proportion of tests with negative side effects

$$H_0: p = 0.005$$

$$H_1: p < 0.005$$

Assume  $H_0$  true.  $X \sim B(2500, 0.005)$

5% significance level

Reject  $H_0$  if  $P(X \leq 3) < 0.05$

Test  $P(X \leq 3) = 0.001525 \dots < 0.05$

The result is significant.

Sufficient evidence to reject  $H_0$

Sufficient evidence to support the medical team's claim

## Worked example

A medical team are testing the negative side effects of a new drug.

They claim that the drug gives negative side effects less than 0.2% of the time.

They test the drug on 4500 patients.

The drug has negative side effects in 5 patients.

Is there enough evidence, at the 1% significance level, to support the medical team's claim?

## Your turn

A medical team are testing the negative side effects of a new drug.

They claim that the drug gives negative side effects less than 0.5% of the time.

They test the drug on 2500 patients.

The drug has negative side effects in 7 patients.

Is there enough evidence, at the 5% significance level, to support the medical team's claim?

$X$  = number of tests with negative side effects

$p$  = probability/proportion of tests with negative side effects

$H_0: p = 0.005$

$H_1: p < 0.005$

Assume  $H_0$  true.  $X \sim B(2500, 0.005)$

5% significance level

Reject  $H_0$  if  $P(X \leq 7) < 0.05$

Test  $P(X \leq 7) = 0.0693 \dots > 0.05$

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to support the medical team's claim

## Worked example

A medical team are testing the negative side effects of a new drug.

They claim that the drug gives negative side effects less than 0.2% of the time.

They test the drug on 4500 patients.

- Find the critical region for this test at the 2% significance level.
- The drug has negative side effects in 5 cases. Comment on this observation in light of the critical region.

## Your turn

A medical team are testing the negative side effects of a new drug.

They claim that the drug gives negative side effects less than 0.5% of the time.

They test the drug on 2500 patients.

- Find the critical region for this test at the 2% significance level.
- The drug has negative side effects in 7 cases. Comment on this observation in light of the critical region.

a)  $X$  = number of tests with negative side effects  
 $p$  = probability/proportion of tests with negative side effects

$$H_0: p = 0.005$$

$$H_1: p < 0.005$$

Under  $H_0$ ,  $X \sim B(2500, 0.005)$

2% significance level

Reject  $H_0$  if  $P(X \leq x) < 0.02$

$$P(X \leq 6) = 0.0342 \dots > 0.02$$

$$P(X \leq 5) = 0.0146 \dots < 0.02$$

$$\therefore x = 5$$

Critical region: Reject  $H_0$  if  $0 \leq X \leq 5$

b) 7 is not in the critical region.

The result is not significant.

Insufficient evidence to reject  $H_0$

Insufficient evidence to support the medical team's claim