

Mark Scheme (Results)

Summer 2018

Pearson Edexcel International GCSE In Mathematics A (4MA1) Paper 1HR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.edexcel.com (some pearson) the UK's largest awarded by Pearson, the UK's largest awarded by Pearson awarded b

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2018
Publications Code 4MA1_1HR_1806_MS
All the material in this publication is copyright
© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded.
 Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
 - Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Types of mark

M marks: method marks

A marks: accuracy marks

o B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- o cao correct answer only
- ft follow through
- o isw ignore subsequent working
- SC special case
- o oe or equivalent (and appropriate)
- o dep dependent
- o indep independent
- o eeoo each error or omission

No working

If no working is shown then correct answers normally score full marks If no working is shown then incorrect (even though nearly correct) answers score no marks.

With working

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.

Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

· Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

International GCSE Maths 4MA1 1HR

Q	Working	Answer	Mark	Notes
1	20 × 14 (= 280)	460	4	M1
	$\frac{20+16}{2} \times (24-14) (=180)$			M1
	"280" + "180"			M1 (dep) on at least one of the previous M marks
				A1
				Total 4 marks
	Alternative scheme 1			
	$(24+14)\div 2 (=19)$ and $(20-16)\div 2 (=2)$	460	4	M1
	$2 \times 19 = 38$) and $16 \times 24 = 384$)			M1
	"38" + "38" + "384"			M1 (dep) on at least one of the previous M marks
				A1
				Total 4 marks
	Alternative scheme 2			
	20 × 24 (= 480)	460	4	M1
	$(20-16) \div 2 (=2)$ and $24-14 (=10)$ $2 \times 10 = 20$			M1
	"480" – "20"			M1 (dep) on at least one of the previous M marks
				A1
				Total 4 marks

2	$1 \times 5 + 3 \times 9 + 5 \times 24 + 7 \times 40 + 9 \times 7 (= 495)$ or $5 + 27 + 120 + 280 + 63 (= 495)$	5.8	4	M2	for at least 4 correct products added (need not be evaluated) If not M2 then award M1 for consistent use of value within interval (including end points) for at least 4 products which must be added OR correct mid-points used for at least 4 products and not added
	"495" ÷ 85			M1	dep on at least M1 Allow division by their $\sum f$ provided addition or total under column seen for $5.8 - 5.824$
					Total 4 marks

3	(a)	Correct R	2	B2	fully correct
		(5,6), (3,6), (3,5)			If not B2 then B1 for correct
					orientation of R but in wrong position
	(b)	Correct T	1	B1	
		(2,-1), (2,-3), (1,-3)			
					Total 3 marks

4	$675 \div (5+4) \times 5 (=375)$	225	3	M1		$M2 675 \div (5+4) \times 3$
	$"375" \div 5 \times 3$			M1	dep M1	
				A1		
						Total 3 marks

5	For example,		No + reason	2	M1	for evaluating E correctly for any
	n	E				value of <i>n</i>
	1	7				
	2	11				
	3	17				
	4	25				
	5	35				
					A1	for No with E evaluated correctly as
						a non-prime number
						Total 2 marks

6	Angle $EBG = 180 - 2 \times 65 (= 50)$ or	27	3	M1	
	Angle $ABE = 180 - (38 + 65) (= 77)$				
	Angle $ABE = 180 - (38 + 65) (= 77)$ and			M1	for a complete method to find
	Angle $ABG = "77" - "50"$				angle <i>ABG</i>
				A1	
					Total 3 marks
	Alternative scheme 1				
	Angle $EBG = 180 - 2 \times 65 (= 50)$ or	27	3	M1	
	Angle $EBC = 103$				
	Angle $EBC = 103$ and			M1	for a complete method to find
	Angle $ABG = 180 - (103 + "50")$				angle ABG
				A1	
					Total 3 marks

7 (a)	4n + 2	2	M1	for $4n + k$ (k may be 0 or absent) oe
			A1	oe
				e.g $6 + (n-1)4$
(b)	4n + 6	1	B1	oe ft part (a) providing M1 in part (a)
				is awarded
				e.g $4(n+1)+2$
				Total 3 marks

8 (a)		1.39×10^{6}	1	B1
(b)		5×10^{-3}	1	B1
				Total 2 marks

9	2.5 - 0.6 = 1.9	2 hours 51 minutes	4	M1	
	3 × 12 × "1.9" (= 68.4)			M1	for using length \times width \times height to find a volume
	"68.4" × 1000 ÷ 400 (= 171 minutes)			M1	for their volume \times 1000 \div 400
				A1	
					Total 4 marks
	Alternative scheme				
	250 - 60 = 190	2 hours 51 minutes	4	M1	
	$300 \times 1200 \times "190" (= 6.84 \times 10^7)$			M1	for using length \times width \times height to find a volume
	"6.84 × 10"" ÷ 10 ⁶ × 1000 ÷ 400 (= 171 minutes)			M1	for their volume $\div 10^6 \times 1000 \div 400$
				A1	
					Total 4 marks

10	16x = 32 or $32y = 144$	(2, 4.5)	3	M1	for a correct sequence of operations
					which leads to 1 equation in one
					unknown, allowing one arithmetical
					error
	$3 \times '2' + 2y = 15 \text{ or } 3x + 2 \times '4.5' = 15$			M1	(dep) substitute found value of one
					variable in one equation
				A1	
					Total 3 marks

11	$72 \times 1000 = 72000 \text{ or } 72 \div 60 = 1.2 \text{ or}$ $72 \div 60 \div 60 = 0.02 \text{ or } 60 \div 60 \times 1000 = 3.6 \text{ or}$	20	3	M1 for at least one of \times 1000 or \div 60
	$\frac{72}{60\times60}\times1000$			M1 (dep) for a complete method
				A1
				Total 3 marks

12 (a)	$6 \times 25 + 6 \times 45 \ (= 150 + 270 = 420)$	20	4	M1	for 6×25 (=150) or 6×45 (=270)
	"150" + "270" – 350 (= 70) or "420" – 350			M1	
	$\frac{"70"}{350} \times 100$			M1	(dep on M2)
				A1	
	Alternative scheme				
	$6 \times 25 + 6 \times 45 \ (= 150 + 270 = 420)$	20	4	M1	for 6×25 (=150) or 6×45 (=270)
	$\frac{"420"}{350} \times 100 = 120$			M1	
	"120" – 100			M1	(dep on M2)
				A1	
(b)	500 000 ÷ 8 (=62 500)	6 250 000	3	M1	
	500 000 ÷8×100			M1	for a complete method
				A1	
					Total 7 marks

13	$\frac{1}{3} + \frac{1}{5} = \frac{8}{15}$ or 0.53 or 53.3% or 53%	900	4	M1
	$1 - \frac{8}{15} = \frac{7}{15}$ or 0.46 or 0.47 or 46.6% or			M1
	47%			
	$420 \div "\frac{7}{15}" (= 900)$ oe			M1
				A1
				Total 4 marks

14	(a)		$8e^{6}f^{9}$	2.	B2	
17	(a)		J Se j	2	B1	for 2 compat tamps in a product of 2
					ы	for 2 correct terms in a product of 3
						terms
	(b)	$3x^2 + 9xy - 4yx - 12y^2$	$3x^2 + 5xy - 12y^2$	2	M1	M1 for 3 correct terms out of 4
						or for 4 correct terms ignoring signs
						or for $3x^2 + 5xy + c$ for any non
						zero value of c
						or for $d + 5xy - 12y^2$ for any non
						zero value of d
					A1	
	(a)	1 2	7	2		for one connect store
	(c)	$\frac{1}{a^2}$ $\frac{3}{a^2}$ $a = a^3$	<u>/</u>	2	M1	for one correct step
		$a^{\frac{1}{2}} \times a = a^{\frac{3}{2}} \text{ or } \frac{a}{a^{-2}} = a^3$	$\frac{1}{2}$			
		$\frac{1}{a^2}$ $\frac{5}{2}$				
		or $\frac{a^{\frac{1}{2}}}{a^{-2}} = a^{\frac{5}{2}}$				
		a^{-2}				
					A1	oe
	(d)	$2^{n}-1$	1	2	M1	for $(2^n - 1)(2^n + 1)$
	. ,					`
		$(2^n-1)(2^n+1)$	$2^{n} + 1$			
					A1	
						Total 8 marks

15	(a)		$\frac{9}{20}$ on first red branch	3	B1	
			20			
			Correct binary structure		B1	
		$\frac{9}{20}$, $\frac{7}{16}$, $\frac{9}{20}$, $\frac{7}{16}$	Labels and correct probabilities on all second branches		B1	
	(b)	$\frac{9'}{20} \times \frac{7'}{16}$	$\frac{63}{320}$ or 0.196(875)	2	M1	
					A1	oe ft diagram Accept 0.20 or better
	(c)	$\frac{9}{20} \times \frac{7}{16} + \frac{11}{20} \times \frac{9}{16}$	$\frac{162}{320} \text{ or } 0.506(25)$	3	M1	for $\frac{11}{20} \times \frac{9}{16}$
					M1	for $\frac{'9'}{20} \times \frac{'7'}{16} + \frac{11}{20} \times \frac{9}{16}$
					A1	oe Accept 0.51 or better
						Total 8 marks

16 (a)	x(x+4) = 12(12+x)	Shown	3	M1
	$x^2 + 4x = 144 + 12x$			M1 for at least one correct expression
				A1 for completion
(b)	$x = \frac{8 \pm \sqrt{(-8)^2 - 4 \times 1 \times (-144)}}{2}$ or	20.6	4	M1 M1 for correctly substituting into the quadratic formula condone one sign error in substitution; allow partial correct evaluation
	$\frac{8\pm\sqrt{8^2-4\times1\times(-144)}}{2} \text{ or } \frac{8\pm\sqrt{-8^2-4\times1\times(-144)}}{2}$ $\frac{8\pm\sqrt{640}}{2} \text{ or } \frac{8\pm\sqrt{(-8)^2576}}{2\times1}$ or $\frac{8\pm8\sqrt{10}}{2}$ NB denominator must be 2×1 or 2 and there must be evidence for correct order of operations in the numerator Allow + instead of ± in the formula			M1 If the first M1 is awarded and an answer of 16.6 or $4 + 4\sqrt{10}$ seen award this M mark
				A1 (dep on M1) 16.6
				B1 (dep on M1) 20.6 - 20.65 ft
				Total 7 marks
17	FDs are 2, 3, 2.8, 0.7, 0.8	Correct histogram	3	M1 for any two correct FD calculations (can be implied by at least two correct bars)
				M1 for any three correct FDs (can be implied by at least three correct bars)
				A1 fully correct histogram
				(SC: B2 for all five bars of correct width with heights in the correct ratio)
				(SC:B1 for three bars of correct width with heights in the correct ratio)
				Total 3 marks

18	$SQ^2 = 8^2 + 12^2 - 2 \times 8 \times 12 \times \cos 120^\circ$	91.4	6		If this mark is awarded then ft on the remaining M marks
	$(SQ) = \sqrt{304}$				for correct order of operations e.g. 64 + 144 + 96 or 304 or 17.4 or $4\sqrt{19}$
	$\frac{\sin R}{\sqrt[n]{304}} = \frac{\sin 27^{\circ}}{9}$			M1	
	$R = \sin^{-1}\left(\frac{\sin 27^{\circ} \times \sqrt{304}}{9}\right)$			M1	can be implied by 61.5833
	61.58			A1	for 61.58 - 61.6
				B1	ft dep M3 180 – "61.6" – 27
		_			Total 6 marks

19	$\frac{\mathrm{d}y}{\mathrm{d}x} = 3x^2 - 27$	108	N	for at least one of $3x^2$ or 27
	$3x^2 - 27 = 0$		N	(dep) for a 2 or 3 term quadratic =0
	$x = \pm 3$		A	1
	When $x = -3$, $b = (-3)^3 - 27(-3) + k (= 54 + k)$ When $x = 3$, $d = 3^3 - 27(3) + k (= -54 + k)$		N	for either substituting $x = 3$ or $x = -3$ into the y expression. Only award this mark if k or a number representing k is in the expression for b or d
	b - d = 54 + k - (-54 + k)			dep on all previous M marks Expressions for b and d must have k or the same number representing k
			F	
				Total 6 marks

20	$h = f\left(\frac{x+1}{2}\right) = 1 + \frac{1}{\frac{x+1}{2}} \left(= 1 + \frac{2}{x+1}\right)$	$\frac{2}{x-1} - 1$ or $\frac{3-x}{x-1}$	4	M1	for $1 + \frac{1}{\frac{x+1}{2}}$
	$\left(y = 1 + \frac{2}{x+1}\right)$ $y - 1 = \frac{2}{x+1} \text{ or } y(x+1) = 1(x+1) + 2$			M1	(dep on M1) for a correct first step to change the subject
	$x + 1 = \frac{2}{y - 1}$ or $xy - x = 3 - y$			M1	(dep on M1)
	$x + 1 = \frac{2}{y - 1} \text{ or } xy - x = 3 - y$ $x = \frac{2}{y - 1} - 1 \text{ or } x = \frac{3 - y}{y - 1}$			A1	oe
					Total 4 marks
	Alternative scheme				
	$h = f\left(\frac{x+1}{2}\right) = 1 + \frac{1}{\frac{x+1}{2}} \left(= 1 + \frac{2}{x+1} = \frac{x+3}{x+1}\right)$	$\frac{3-x}{x-1}$	4	M1	for $1 + \frac{1}{\frac{x+1}{2}}$
				M1	(dep on M1) for a correct first step to change the subject
	xy - x = 3 - y			M1	(dep on M1)
	$x = \frac{3-y}{y-1}$			A1	oe
					Total 4 marks
Note: Allow can	didates to swap x and y when finding the inverse		1	I	

21 (a)	$x(x^2-1)$ or $(x^2-x)(x+1)$	x^3-x	1	B1 for correct expansion of a pair of
				brackets and then $x^3 - x$ written
				down
(b)	(One of the numbers) is even or multiple of 2 or 2 is a factor	Proof	3	M1
	(One of the numbers) is a multiple of 3 or 3 is a factor			M1
	Hence a multiple of 6			A1
				Total 4 marks

22	$\frac{4}{3} \times \pi \times R^3 - \frac{4}{3} \times \pi \times 1.2^3$ or	3.9	5	M1	for an expression for the volume of the inner sphere
	$\frac{4}{3} \times \pi \times (1.2 + t)^3 - \frac{4}{3} \times \pi \times 1.2^3$				
	$\frac{\frac{4}{3} \times \pi \times (1.2 + t)^3 - \frac{4}{3} \times \pi \times 1.2^3}{\left(\frac{4\pi}{3}R^3 - \frac{4}{3} \times \pi \times 1.2^3\right) \times 2700 = 1980}$			M1	
	$\frac{4\pi}{3}R^3 = \frac{4}{3} \times \pi \times 1.2^3 + \frac{1980}{2700}$ $= 7.238229474 + 0.73333333$ $= 7.97(1562807)$			M1	for a correct expression or sight of 7.23(8229474) + 0.73(33333) or sight of 7.97(1562807)
	$R = \sqrt[3]{\left(\frac{3}{4\pi} \times (\frac{4}{3} \times \pi \times 1.2^3 + \frac{1980}{2700})\right)} = 1.2392$ $1.2392 - 1.2 = 0.0392$			M1	for a correct expression or sight of $\sqrt[3]{1.90(3070437)}$ or sight of 1.23(9229151) or sight of 0.0392(29151)
				A1	for 3.9 – 3.92
					Total 5 marks
23	(First term = 3 and last term = 999) or $a = 3$ and $d = 3$	166 833	4	M1	
	999 ÷ 3 (= 333)			M1	for finding the number of terms Allow $1000 \div 3 = 333.3 = 333$
	Sum = $\frac{333}{2}$ (3+999) or Sum = $\frac{333}{2}$ (2×3+(333-1)3)			M1	for using a correct method to find the sum
	Sum = $\frac{333}{2} (2 \times 3 + (333 - 1)3)$				

A1

Total 4 marks





