



---

# LEVEL 2 CERTIFICATE

# Further Mathematics

Paper 2 8360/2 Calculator

Mark scheme

---

8360

June 2017

---

Version: 1.0 Final

---

---

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Assessment Writer.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

---

## Glossary for Mark Schemes

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

<b>M</b>	Method marks are awarded for a correct method which could lead to a correct answer.
<b>A</b>	Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied.
<b>B</b>	Marks awarded independent of method.
<b>ft</b>	Follow through marks. Marks awarded for correct working following a mistake in an earlier step.
<b>SC</b>	Special case. Marks awarded for a common misinterpretation which has some mathematical worth.
<b>M dep</b>	A method mark dependent on a previous method mark being awarded.
<b>B dep</b>	A mark that can only be awarded if a previous independent mark has been awarded.
<b>oe</b>	Or equivalent. Accept answers that are equivalent. eg accept 0.5 as well as $\frac{1}{2}$
<b>[a, b]</b>	Accept values between a and b inclusive.
<b>[a, b)</b>	Accept values $a \leq \text{value} < b$
<b>3.14...</b>	Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416
<b>Use of brackets</b>	It is not necessary to see the bracketed work to award the marks.

Examiners should consistently apply the following principles

### **Diagrams**

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

### **Responses which appear to come from incorrect methods**

Whenever there is doubt as to whether a student has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the student. In cases where there is no doubt that the answer has come from incorrect working then the student should be penalised.

### **Questions which ask students to show working**

Instructions on marking will be given but usually marks are not awarded to students who show no working.

### **Questions which do not ask students to show working**

As a general principle, a correct response is awarded full marks.

### **Misread or miscopy**

Students often copy values from a question incorrectly. If the examiner thinks that the student has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

### **Further work**

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

### **Choice**

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

### **Work not replaced**

Erased or crossed out work that is still legible should be marked.

### **Work replaced**

Erased or crossed out work that has been replaced is not awarded marks.

### **Premature approximation**

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

### **Continental notation**

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the student intended it to be a decimal point.

Q	Answer	Mark	Comments
1(a)	$\frac{3-5 \times 20}{2}$ or $\frac{3-100}{2}$ or $(-)\frac{97}{2}$ or $(-)\text{48.5}$ or $\frac{3-5 \times 8}{2}$ or $\frac{3-40}{2}$ or $(-)\frac{37}{2}$ or $(-)\text{18.5}$ or $12 \times (-)\frac{5}{2}$	M1	oe
	(-)30	A1	Accept if both 30 and -30 are seen
	<b>Additional Guidance</b>		
1(b)	$-\frac{3}{2}$ or $-1\frac{1}{2}$ or $-1.5$	B1	oe
	<b>Additional Guidance</b>		
	Condone $\frac{3}{-2}$ or $n \rightarrow -1.5$ or $-1\frac{1}{2} \rightarrow \infty$		B1
	$-\frac{3n}{2n}$		B1
	$-\frac{3n}{2n}$ not processed		B0
	$\frac{3}{0-2}$ not processed		B0
$-1.5n$		B0	

<b>2(a)</b>	$\begin{pmatrix} 13 & -2 \\ 6 & 1 \end{pmatrix}$	B2	B1 13 or -2 or 6 or 1 in correct position in a 2 by 2 matrix
	<b>Additional Guidance</b>		
	Condone missing brackets for B2 or B1 if numbers in a 2 by 2 array		
	Brackets may be square or curly etc		
	Ignore commas and fraction lines		
	$\begin{pmatrix} 13 & -2 \\ 6 & 1 \end{pmatrix}$ followed by further work		B1
<b>2(b)</b>	$5k = 11 - 3k$ or $2k = 11 - 6k$ or $11 - 3k = \frac{5}{2}(11 - 6k)$ or $8k = 11$	M1	oe Any one correct equation
	$\frac{11}{8}$ or $1\frac{3}{8}$ or 1.375 with no incorrect equation seen	A1	oe
	<b>Additional Guidance</b>		
	$\begin{pmatrix} 5k \\ 2k \end{pmatrix} = \begin{pmatrix} 11-3k \\ 11-6k \end{pmatrix}$ with no further correct work		M0
	Ignore subsequent attempt to convert $\frac{11}{8}$ to a mixed fraction or decimal		M1A1
	Ignore subsequent attempt to convert $1\frac{3}{8}$ to an improper fraction or decimal		M1A1
	Ignore subsequent rounding or truncation of 1.375		M1A1
	Answer only 1.37 or 1.38 or 1.4		M0
T & I is 2 or zero			

<b>2(c)</b>	Valid explanation	B1	eg Number of columns of <b>B</b> does not equal number of rows of <b>A</b>
	<b>Additional Guidance</b>		
	'2 by 1' (or '2 × 1') matrix means 2 rows and 1 column		
	'First matrix' means B and 'second matrix' means A		
	B columns $\neq$ A rows		B1
	B rows $\neq$ A columns		B0
	B columns $\neq$ A columns		B0
	B rows $\neq$ A rows		B0
	Number of rows in second matrix cannot be more than number of columns in first matrix		B1
	B has 1 column, A has 2 rows		B1
	A should only have 1 row		B1
	A has too many rows		B1
	B only has one column		B1
	B needs another column		B1
	It is a 2 x 1 multiplied by a 2 x 2		B1
	There's nothing to multiply the 3 by		B1
	It is a 2 x 2 multiplied by 2 x 1		B0
	It is a 1 by 2 multiplied by a 2 by 2		B0
	B values can't multiply with all the A values		B0
	They are not compatible		B0
	Because the dimensions of A and B are different		B0
	Can't work it out this way round		B0
	Can work out AB but not BA		B0
B has to be a 2 by 2 matrix		B0	

<b>3(a)</b>	3 (×) 455 or 5 (×) 273 or 7 (×) 195 or 13 (×) 105 or 15 (×) 91 or 21 (×) 65 or 35 (×) 39 or 3 (×) 5 (×) 7 (×) 13	M1	oe eg $1365 \div 5 = 273$ Any order Must be integers May be seen in a factor tree or repeated division
	3 5 91 or 3 7 65 or 3 13 35 or 5 7 39 or 5 13 21 or 7 13 15	A1	Any order Must be integers
	<b>Additional Guidance</b>		
	If using division the correct answer must be seen for M1		
	Correct answer can be implied by working lines eg 3 (×) 5 (×) 91 with blank answer line		M1A1
	Answer line correct		M1A1
	Allow inclusion of 1 for M1 eg 1 (×) 3 (×) 455		M1
<b>3(b)</b>	$b(a - 11)$ or $-b(11 - a)$	M1	Implied by square numbers $> 1$ used eg1 $4(36 - 11)$ eg2 $9(16 - 11)$
	$a = 36$ and $b = \text{square number} > 1$ with working for M1 seen	A1	Must be in correct order Allow unprocessed squares eg $a = 6^2$ and $b = 5^2$  SC1 $a = 36$ and $b = \text{square number} > 1$ without working for M1 seen
	<b>Additional Guidance</b>		
	$b(a - 11) = 0$ or $b(a - 11)$ with further work		M1
	Answer line takes precedence over working lines		
Embedded answer eg $81(36 - 11)$		M1A0	



<b>4</b>	$\left(\frac{56}{4}\right)^3$ or $14^3$ or $4^3x = 56^3$ or $64x = 175\,616$ or $\frac{56^3}{x} = 4^3$	M1	oe  oe equation in $x^{(1)}$ or $\frac{1}{x^{(1)}}$
	2744	A1	
	<b>Additional Guidance</b>		
	$\sqrt[3]{x} = \frac{56}{4}$ or $\sqrt[3]{x} = 14$ with no correct further work		M0
	$56x^{-\frac{1}{3}} = 4$		M0
	Solving $\frac{56}{3x} = 4$		M0
	Answer $14^3$ with 2744 not seen in working		M1A0
Embedded solution		M1A0	

<b>5</b>	<b>Alternative method 1</b>		
	$\frac{a+4}{2} = 3a$ or $3a - a = 4 - 3a$ or $a + \frac{4-a}{2} = 3a$ or $4 - \frac{4-a}{2} = 3a$ or $4 - a = 2(3a - a)$	M1	oe
	$6a - a = 4$ or $3a - a + 3a = 4$ or $2a - a - 6a = -4$ or $8 - 4 = 6a - a$ or $4 = 4a + a$ or $5a = 4$	M1dep	oe Allow eg $3a \times 2$ for $6a$ Terms collected
	$\frac{4}{5}$ or 0.8	A1	oe
	<b>Alternative method 2</b>		
	$\frac{8-6}{3a-a} = \frac{10-6}{4-a}$ or $\frac{8-6}{3a-a} = \frac{10-8}{4-3a}$ or $\frac{10-6}{4-a} = \frac{10-8}{4-3a}$	M1	oe eg fractions inverted
	$8a + 2a = 8$ or $6a + 4a = 8$ or $-12a + 2a = 8 - 16$ or $5a = 4$	M1dep	oe Allow eg $2a \times 4$ for $8a$ Terms collected
	$\frac{4}{5}$ or 0.8	A1	oe

**Alternative method 3 and Additional Guidance continue on the next page**

<b>5 cont</b>	<b>Alternative method 3</b>		
	$(8 - 6)^2 + (3a - a)^2$ $= (10 - 8)^2 + (4 - 3a)^2$ or $5a^2 - 24a + 16$ (= 0) or $(10 - 6)^2 + (4 - a)^2$ $= 2^2((8 - 6)^2 + (3a - a)^2)$ or $15a^2 + 8a - 16$ (= 0) or $(10 - 6)^2 + (4 - a)^2$ $= 2^2((10 - 8)^2 + (4 - 3a)^2)$ or $35a^2 - 88a + 48$ (= 0)	M1	oe Using $PM^2 = MQ^2$ or $PQ^2 = 4PM^2$ or $PQ^2 = 4MQ^2$
	$(5a - 4)(a - 4)$ (= 0) or $\frac{-24 \pm \sqrt{(-24)^2 - 4 \times 5 \times 16}}{2 \times 5}$ or $(5a - 4)(3a + 4)$ (= 0) or $\frac{-8 \pm \sqrt{8^2 - 4 \times 15 \times -16}}{2 \times 15}$ or $(5a - 4)(7a - 12)$ (= 0) or $\frac{-88 \pm \sqrt{(-88)^2 - 4 \times 35 \times 48}}{2 \times 35}$	M1dep	oe eg $\frac{12}{5} \pm \sqrt{\frac{64}{25}}$ or $-\frac{4}{15} \pm \sqrt{\frac{256}{225}}$ or $\frac{44}{35} \pm \sqrt{\frac{256}{1225}}$ Correct attempt to solve their 3-term quadratic Allow recovery of brackets in formula Allow eg $24^2$ for $(-24)^2$ Implied by correct solutions to their 3-term quadratic seen
	$\frac{4}{5}$ or 0.8	A1	oe Must reject solution 4 or $-\frac{4}{3}$ or $\frac{12}{7}$
	<b>Additional Guidance</b>		
	Terms must be collected but do not have to be processed for M1dep eg (Alt 1) $a + 4 = 6a$ needs terms collecting to $4 = 6a - a$ for M1dep		
	Rejection of solution is implied by only $\frac{4}{5}$ on answer line		

	$\sin 38 = \frac{r}{20}$ or $\cos (90 - 38) = \frac{r}{20}$ or $\frac{r}{\sin 38} = \frac{20}{\sin 90}$ or $\frac{\sin 38}{r} = \frac{\sin 90}{20}$	M1	oe Any letter
	$20 \times \sin 38$ or $20 \times \cos (90 - 38)$ or $\frac{20}{\sin 90} \times \sin 38$	M1dep	oe M2 $\sqrt{20^2 - (20 \cos 38)^2}$ M2 $\frac{\sqrt{20^2 + 20^2 - 2 \times 20 \times 20 \times \cos(38 \times 2)}}{2}$
	12.3(...)	A1	SC2 Angle VAC = 38 seen on diagram <b>and</b> answer 15.76(...) or 15.8
<b>Additional Guidance</b>			
<b>6</b>	If trigonometry and Pythagoras are used it must be a fully correct method that would lead to the correct value of $r$ for M2		
	If cosine rule with angle $(38 \times 2)$ used it must be a fully correct method that would lead to the correct value of $r$ for M2		
	Answer 15.76(...) or 15.8 but angle VAC = 38 not seen		Zero
	12.3(...) seen and angle 38 in correct place with further work eg $20 \sin 38 = 12.3$ $\sqrt{20^2 - 12.3^2} = 15.8$		Zero
	$\sin 38 \times 20$ (even if subsequently evaluates $\sin 760$ )		M2
	Throughout, accept opp or o for $r$ eg $\sin 38 = \frac{\text{opp}}{20}$		M1
	$\sin = \frac{r}{20}$ or $\sin \theta = \frac{r}{20}$ (unless recovered)		M0
	Answer 12.3(...) coming from scale drawing		M2A1
	Answer 12 coming from scale drawing		Zero
	12.3(...) seen with no further work followed by answer 12		M2A1

<b>7</b>	<b>Alternative method 1</b>		
	( $x$ -coordinate of $A =$ ) 10 <b>and</b> ( $y$ -coordinate of $B =$ ) 8	B1	May be implied on diagram eg 10 written next to $A$ <b>and</b> 8 written next to $B$
	( $x$ -coordinate of $P =$ ) $\frac{2}{2+3} \times$ their 10 or $\frac{2 \times \text{their } 10 + 3 \times 0}{2+3}$ or 4	M1	oe their 10 must be their $x$ -coordinate of $A$ May be seen on diagram
	(area of triangle $OBP =$ ) $\frac{1}{2} \times$ their 8 $\times$ their 4	M1dep	oe their 8 must be their $y$ -coordinate of $B$
	16	A1ft	ft B0M2
	<b>Alternative method 2</b>		
	( $x$ -coordinate of $A =$ ) 10 <b>and</b> ( $y$ -coordinate of $B =$ ) 8	B1	May be implied on diagram eg 10 written next to $A$ <b>and</b> 8 written next to $B$
	(area of triangle $OAB =$ ) $\frac{1}{2} \times$ their 10 $\times$ their 8 or 40	M1	oe
	(area of triangle $OBP =$ ) $\frac{2}{2+3} \times$ their 40	M1dep	oe eg their $40 - \frac{3}{2+3} \times$ their 40
	16	A1ft	ft B0M2

**Alternative methods 3 and 4 and Additional Guidance continue on the next two pages**

<b>7</b> <b>cont</b>	<b>Alternative method 3</b>		
	( $x$ -coordinate of $A =$ ) 10 <b>and</b> ( $y$ -coordinate of $B =$ ) 8	B1	May be implied on diagram eg 10 written next to $A$ <b>and</b> 8 written next to $B$
	(area of triangle $OAB =$ ) $\frac{1}{2} \times$ their 10 $\times$ their 8 or 40	M1	oe
	( $y$ -coordinate of $P =$ ) $\frac{3}{2+3} \times$ their 8 or 4.8 <b>and</b> (area of triangle $OPA =$ ) $\frac{1}{2} \times$ their 10 $\times$ their 4.8 or 24 <b>and</b> (area of triangle $OBP =$ ) their 40 – their 24	M1dep	oe their 8 must be their $y$ -coordinate of $B$ $y$ -coordinate of $P$ may be seen on diagram
	16	A1ft	ft BOM2

**Alternative method 4 and Additional Guidance continue on the next page**

<b>7 cont</b>	<b>Alternative method 4</b>		
	( $x$ -coordinate of $A =$ ) 10 <b>and</b> ( $y$ -coordinate of $B =$ ) 8	B1	May be implied on diagram eg 10 written next to $A$ <b>and</b> 8 written next to $B$
	( $AB =$ ) $\sqrt{\text{their } 10^2 + \text{their } 8^2}$ or $\sqrt{100 + 64}$ or $\sqrt{164}$ or $2\sqrt{41}$ or 12.8(...) <b>and</b> ( $BP =$ ) $\frac{2}{2+3} \times \text{their } 12.8(\dots)$ or 5.12(...) <b>and</b> (angle $OBP =$ ) $\tan^{-1} \frac{\text{their } 10}{\text{their } 8}$ or 51.3(...)	M1	oe  their 10 must be their $x$ -coordinate of $A$ their 8 must be their $y$ -coordinate of $B$
	(area of triangle $OBP =$ ) $\frac{1}{2} \times \text{their } 8 \times \text{their } 5.12$ $\times \sin \text{ their } 51.3$	M1dep	oe their 8 must be their $y$ -coordinate of $B$
	16	A1ft	ft B0M2
	<b>Additional Guidance</b>		
	$A = 10$ and $B = 8$		B1
	$A(8, 0)$ and $B(0, 10)$ is B0 but can subsequently score up to M2A1ft (answer 16)		
	$A(0, 10)$ and $B(8, 0)$ is B0 but can score up to M2A1ft if uses $x$ -coordinate of $A$ as 10 and $y$ -coordinate of $B$ as 8 (answer 16)		
	$A(0, 8)$ and $B(10, 0)$ is B0 but can score up to M2A1ft if uses $x$ -coordinate of $A$ as 8 and $y$ -coordinate of $B$ as 10 (answer 16)		
Area triangle $OBP$ may be seen as the sum of two right-angled triangles			
Area triangle $OBP$ may be seen as area trapezium $OBPX$ – area triangle $OPX$ $X$ is on the $x$ -axis with $PX$ perpendicular to the $x$ -axis			
Allow marks for valid working seen even if not subsequently used			
15.9(...) $\rightarrow$ answer 16 Answer 15.9(...)		4 marks B1M2A0	

<b>8</b>	<b>Alternative method 1</b>		
	$(BC =) 12$	B1	Allow as two 6s labelled on $BC$ after perpendicular drawn from $A$
	their $12^2 = 7^2 + 8^2$ $- 2 \times 7 \times 8 \times \cos A$ or $144 = 49 + 64 - 112 \cos A$ or $144 = 113 - 112 \cos A$ or $\frac{7^2 + 8^2 - \text{their } 12^2}{2 \times 7 \times 8}$ or $\frac{49 + 64 - 144}{112}$ or $-\frac{31}{112}$ or $[-0.277, -0.27]$ or $-0.28$	M1	oe Do not allow if their 12 comes from use of Pythagoras' theorem ie $(BC =) \sqrt{7^2 + 8^2}$ or $\sqrt{113}$ or $10.6(\dots)$ is B0M0
	$\cos^{-1} \left( \frac{7^2 + 8^2 - \text{their } 12^2}{2 \times 7 \times 8} \right)$	M1dep	oe May be implied by final answer
	[106, 106.1]	A1ft	Only ft B0M2
	<b>Alternative method 2</b>		
	$(BC =) 12$	B1	Allow as two 6s labelled on $BC$ after perpendicular drawn from $A$
	(angle $ABC =$ ) $\cos^{-1} \left( \frac{7^2 + \text{their } 12^2 - 8^2}{2 \times 7 \times \text{their } 12} \right)$ or 39.8... <b>and</b> $\sin A = \frac{\sin \text{their } 39.8}{8} \times \text{their } 12$ or $\sin A = 0.96\dots$	M1	oe eg works out angle $ACB$ ( $= 34.09(\dots)$ or $34.1$ ) and uses sine rule Do not allow if their 12 comes from use of Pythagoras' theorem ie $(BC =) \sqrt{7^2 + 8^2}$ or $\sqrt{113}$ or $10.6(\dots)$ is B0M0
	$180 - \sin^{-1}(\text{their } 0.96\dots)$	M1dep	oe May be implied by final answer
	[106, 106.1]	A1ft	Only ft B0M2

**Additional Guidance continues on the next page**



	<b>Additional Guidance</b>	
<b>8 cont</b>	$\cos^{-1}$ or $\cos^{-1}$ ans does not score M1dep unless recovered	
	For the M1dep must have correct rearrangement but allow arithmetic errors	
	Answer outside range is A0 eg 106.2(...) from $\cos^{-1}(-0.28)$	

<b>9</b>	<b>Alternative method 1</b>		
	$-\frac{11}{5} < x \leq \frac{5}{5}$ or $-2.2 < x \leq \frac{5}{5}$	M1	oe eg $x \leq \frac{5}{5}$ and $x > -\frac{11}{5}$
	$-\frac{11}{5} < x \leq 1$ or $-2.2 < x \leq 1$ or $-2 \leq x \leq 1$ or $-2, -1, 0, 1$	A1	oe eg $x \leq 1$ and $x > -\frac{11}{5}$
	$6x - 4x \leq 4 - 7$ or $2x \leq -3$	M1	oe Collects terms
	$x \leq -\frac{3}{2}$ or $x \leq -1.5$ or $x < -\frac{3}{2}$ or $x < -1.5$ or $x \leq -2$ or $-2, -3$ ( $, -4, \dots$ )	A1	$-2.2 < x \leq -1.5$ or $-2 \leq x \leq -1.5$ implies M1A1M1A1
	$-2$ with no other values given	A1	Must have gained M1A1M1A1
	<b>Alternative method 2</b>		
	Shows that $-2$ satisfies either $-11 < 5x \leq 5$ or $6x + 7 \leq 4x + 4$	M1	eg $-11 < -10 \leq 5$ or $5x = -10$ and yes
	Shows that $-2$ satisfies both $-11 < 5x \leq 5$ and $6x + 7 \leq 4x + 4$	A1	
	Shows that $-1$ does not satisfy $6x + 7 \leq 4x + 4$ <b>or</b> shows that $-3$ does not satisfy $-11 < 5x \leq 5$	M1	eg $-6 + 7 > -4 + 4$
	Shows that $-1$ does not satisfy $6x + 7 \leq 4x + 4$ <b>and</b> shows that $-3$ does not satisfy $-11 < 5x \leq 5$	A1	
	$-2$ with no other values given	A1	Must have gained M1A1M1A1


**Alternative methods 3 and 4 and Additional Guidance continue on the next two pages**

<b>Alternative method 3</b>			
<b>9</b>	$-\frac{11}{5} < x \leq \frac{5}{5}$ or $-2.2 < x \leq \frac{5}{5}$	M1	oe eg $x \leq \frac{5}{5}$ and $x > -\frac{11}{5}$
	$-\frac{11}{5} < x \leq 1$ or $-2.2 < x \leq 1$ or $-2 \leq x \leq 1$ or $-2, -1, 0, 1$	A1	oe eg $x \leq 1$ and $x > -\frac{11}{5}$
	Shows that $-2$ satisfies $6x + 7 \leq 4x + 4$ <b>or</b> shows that $-1$ does not satisfy $6x + 7 \leq 4x + 4$	M1	eg $6 \times -2 + 7 = -5$ and $4 \times -2 + 4 = -4 \checkmark$
	Shows that $-2$ satisfies $6x + 7 \leq 4x + 4$ <b>and</b> shows that $-1$ does not satisfy $6x + 7 \leq 4x + 4$	A1	
	$-2$ with no other values given	A1	Must have gained M1A1M1A1

**Alternative method 4 and Additional Guidance continue on the next page**

<b>9</b>	<b>Alternative method 4</b>		
	$6x - 4x \leq 4 - 7$ or $2x \leq -3$	M1	oe Collects terms
	$x \leq -\frac{3}{2}$ or $x \leq -1.5$ or $x < -\frac{3}{2}$ or $x < -1.5$ or $x \leq -2$ or $-2, -3$ (, $-4, \dots$ )	A1	
	Shows that $-2$ satisfies $-11 < 5x \leq 5$ <b>or</b> shows that $-3$ does not satisfy $-11 < 5x \leq 5$	M1	eg $-11 < -10 \leq 5$ or $5x = -10$ and yes
	Shows that $-2$ satisfies $-11 < 5x \leq 5$ <b>and</b> shows that $-3$ does not satisfy $-11 < 5x \leq 5$	A1	
	$-2$ with no other values given	A1	Must have gained M1A1M1A1
	<b>Additional Guidance</b>		
	Allow eg max 1 and min $-2.2$ for $-2.2 < x \leq 1$ , unless contradicted by a list of values		
	Condone omission of non-critical values from lists eg $-2, -1, 1$		
	Using = signs when solving inequalities can score M marks only unless recovered		
Incorrect notation eg $\leq$ for $<$ can score M marks only			
If answers to trials evaluated they must be correct			
Choose the scheme that favours the student			
$-2$ identified as the only integer with no valid working		Zero	

<b>10</b>	<b>Alternative method 1</b>		
	$\frac{1}{2} \times x \times x \times \sin 150$ or $\frac{1}{4} x^2$ or $\frac{1}{2} \times b \times c \times \sin 150 = 57.76$ or $\frac{1}{4} \times b \times c = 57.76$	M1	oe Any letter(s)
	$x^2 = \frac{57.76 \times 2}{\sin 150}$ or $x^2 = 57.76 \times 4$ or $x^2 = 231(.04)$ or $\frac{1}{2} x = \sqrt{57.76}$ or $\sqrt{231(.04)}$ or $2\sqrt{57.76}$	M1dep	oe eg $x^2 = \frac{57.76}{\frac{1}{2} \sin 150}$  Must have either $x^2 =$ or $\frac{1}{2} x = \sqrt{57.76}$  or $\sqrt{231(.04)}$ or $2\sqrt{57.76}$  Any letter
	15.2	A1	
	<b>Alternative method 2</b>		
	$\frac{1}{2} \times x \times x \cos \frac{150}{2} \times \sin \frac{150}{2}$  $= \frac{57.76}{2}$	M1	oe Any letter
	$x^2 = \frac{57.76}{\cos \frac{150}{2} \sin \frac{150}{2}}$ or $x^2 = 231(.04)$ or $\sqrt{231(.04)}$ or $2\sqrt{57.76}$	M1dep	oe Must have either $x^2 =$ or $\sqrt{231(.04)}$ or $2\sqrt{57.76}$  Any letter
	15.2	A1	
	<b>Additional Guidance</b>		
	Do not allow 15 as a misread of 150		
$x$ can be $b$ or $AB$ or $AC$ etc			
$b$ and $c$ can be $a$ and $b$ or $AB$ and $AC$ etc			

<b>11(a)</b>	Straight line between $(-2, 7)$ and $(0, 3)$	B1	Tolerance of $\pm 1$ small square Allow line to be extended
	Points $(0, 3)$ $(1, 4)$ $(2, 3)$ $(3, 0)$ $(4, -5)$	M1	Tolerance of $\pm 1$ small square May be plotted or seen in a table Points can be implied
	Correct smooth parabolic curve with maximum at $(1, 4)$	A1	Tolerance of $\pm 1$ small square Allow (ruled) straight line between $(3, 0)$ and $(4, -5)$ Curve passing through all correct points within tolerance scores M1A1
	Straight line between $(4, -5)$ and $(5, 0)$	B1	Tolerance of $\pm 1$ small square Allow line to be extended
	<b>Additional Guidance</b>		
	Ignore extra points plotted		
	Tolerance of $\pm 1$ small square means it is on the edges of or within the shaded area		
			
	Points only can score a maximum of M1		
	Ruled straight lines for curve apart from between $(3, 0)$ and $(4, -5)$		
If all 4 marks would be awarded but either (i) graph has a line or a curve that extends beyond the individual domains or (ii) the curve does not meet a line at a cusp			3 marks

<b>11(b)</b>	$-5 \leq f(x) \leq 7$ or $7 \geq f(x) \geq -5$ or $[-5, 7]$	B2ft	Correct or ft their graph in (a) for B2 ft their graph in (a) for B1 B1ft $-5 \leq f(x)$ or $f(x) \leq 7$ on their own or embedded within an interval for $f(x)$ or only $-5$ and $7$ chosen eg $-5 < f(x) < 7$
	<b>Additional Guidance</b>		
	Allow $f(x)$ to be $y$ or $f$ or $fx$ eg1 $-5 \leq y \leq 7$ eg2 $f \leq 7$		B2 B1
	Allow as two inequalities $f(x) \geq -5$ (and/or) $f(x) \leq 7$		B2
	ft their graph if incomplete eg no graph drawn for $-2 \leq x < 0$ but otherwise correct and answer $-5 \leq f(x) \leq 4$		B2ft
	ft their graph if drawn for $x$ values beyond $[-2, 5]$ eg1 straight line from $(-3, 8)$ to $(6, -1)$ and answer $-1 \leq y \leq 8$ eg2 straight line from $(-3, 8)$ to $(6, -1)$ and answer $f(x) \leq 8$		B2ft B1ft
	Straight line from $(-2, 9)$ to $(6, -7)$ and answer $-7 \leq y \leq 9$		B2ft
	Straight line from $(0, 9)$ to $(5, -4)$ and answer $-4 \leq f(x) \leq 9$		B2ft
	B2ft (or B1ft) can be awarded for a range beyond $[-7, 9]$ if it is clear from working (eg a table of values) where the answer is from		
	$-5$ to $7$ inclusive is B2 whereas $-5$ to $7$ is B1		
	B1 for a correct inequality embedded eg1 $-5 < f(x) \leq 7$ eg2 $-5 \leq f(x) < 0$ eg3 $-2 \leq y \leq 7$		B1 B1 B1
	For B1 ignore incorrect notation if only $-5$ and $7$ chosen eg1 $-5 \leq x \leq 7$ eg2 $-5 < x \leq 7$ eg3 $-5 \geq f(x) \geq 7$ eg4 $-5, 7$		B1 B1 B1 B1
	$\{-5, -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6, 7\}$		B0
	Working out a statistical range eg $-5$ to $7 = 12$		B0

<b>12(a)</b>	$3(25 - x^2)$ or $-3(x^2 - 25)$ or $(15 + 3x)(5 - x)$ or $(x + 5)(15 - 3x)$	M1	oe partial factorisation eg $-(3x + 15)(x - 5)$ Brackets in either order Do not allow $-(3x^2 - 75)$
	$3(5 + x)(5 - x)$ or $3(-x - 5)(x - 5)$ or $-3(x + 5)(x - 5)$ or $-3(5 - x)(-x - 5)$	A1	
	<b>Additional Guidance</b>		
	$(-x + 5)$ is equivalent to $(5 - x)$ etc		
	Do not allow A1 for incorrect notation in final answer eg $(5 + x)3(5 - x)$		M1A0
	Do not allow A1 for use of multiplication signs in final answer eg $3 \times (5 + x) \times (5 - x)$		M1A0
	Correct answer followed by incorrect further work		M1A0



<b>12(b)</b>	<b>Alternative method 1</b>		
	$9n^2 + 3n + 3n + 1$ or $9n^2 + 6n + 1$ or $9n^2 - 3n - 3n + 1$ or $9n^2 - 6n + 1$	M1	oe Terms may be seen in a grid
	$12n$ with no incorrect working	A1	Brackets can be recovered
	<b>Alternative method 2</b>		
	$(3n + 1 + 3n - 1)(3n + 1 - (3n - 1))$ or $(3n + 1 + 3n - 1)(3n + 1 - 3n + 1)$	M1	oe Brackets around $3n - 1$ can be recovered
	$12n$	A1	
	<b>Additional Guidance</b>		
	Alt 1 $12n$ may come from incorrect working eg1 $3n^2 + 6n + 1 - (3n^2 - 6n + 1) = 12n$ eg2 $9n^2 + 3n + 1 - (9n^2 - 3n + 1) = 12n$		M0A0 M0A0
	Alt 1 Recovery of brackets eg1 $9n^2 + 6n + 1 - 9n^2 - 6n + 1 = 12n$ eg2 $9n^2 + 6n + 1 - 9n^2 - 6n + 1 = 2$		M1A1 M1A0
	Alt 2 Recovery of brackets eg1 $(3n + 1 + 3n - 1)(3n + 1 - 3n - 1) = 12n$ eg2 $(3n + 1 + 3n - 1)(3n + 1 - 3n - 1) = 0$		M1A1 M0A0
Do not allow A1 for use of multiplication signs in final answer eg $12 \times n$ with no incorrect working		M1A0	

<b>13</b>	Single correct fraction with terms processed	M1	eg1 $\frac{600a^5 + 1200a^4}{36a^3 + 72a^2}$ eg2 $\frac{50a^3 + 100a^2}{3a + 6}$ Only bracket allowed is $(a + 2)$ eg $\frac{50a^4(a + 2)}{3a^3 + 6a^2}$ (scores M2)
	Factorises correctly using $(a + 2)$	M1	Only needs to be seen once eg1 $\frac{8a}{3a + 6} \times \frac{5(a + 2)}{3a^2} \div \frac{4}{15a^3}$ eg2 $\frac{8a}{3(a + 2)} \times \frac{5a + 10}{3a^2} \times \frac{15a^3}{4}$ Award M2 for fully correct unprocessed expression with full cancelling seen eg $\frac{\cancel{8}^2 a}{3(\cancel{a + 2})} \times \frac{5(\cancel{a + 2})}{\cancel{3} a^2} \times \frac{\cancel{5}^1 \cancel{15}^3 a^3}{\cancel{4}^1}$ or $\frac{2a}{3} \times 5 \times 5a$ oe
	$\frac{50a^2}{3}$ or $16\frac{2}{3}a^2$ or $16.\dot{6}a^2$	A1	
	<b>Additional Guidance</b>		
	$\frac{50 \times a \times a}{3}$		M2A0
	A correct single fraction with $(a + 2)$ cancelled will be M2 eg1 $\frac{250a^2}{15}$ eg2 $\frac{50a^4}{3a^2}$		M2A0
	$\frac{8a}{3} \times \frac{5(a + 2)}{3a^2} \times \frac{15a^3}{4}$		M0M1A0
	$3a + 6 = 3(a + 2)$ with no other valid working		M0M1A0
	Brackets other than $(a + 2)$ may be seen $\frac{10a^2(5a + 10)}{3a + 6}$		M0M0
Correct answer followed by incorrect further work		M2A0	
Allow one miscopy for up to M2A0			

<b>14</b>	<b>Alternative method 1</b>		
	$-\frac{1}{4}$ or $-0.25$	B1	gradient of $x + 4y = 74$ Do not allow embedded May be implied
	(gradient =) $\frac{-1}{\text{their } -\frac{1}{4}}$ or 4	M1	ft their $-\frac{1}{4}$ Only ft a non-zero numerical value Implied by $y = 4x + b$ or $a = 4$ (B1M1)
	$(y =) \frac{74-2}{4}$ or $\frac{72}{4}$ or 18	M1	oe May be seen on diagram
	their $18 = \text{their } 4 \times 2 + b$ or $y - \text{their } 18 = \text{their } 4(x - 2)$	M1dep	oe dep on M2
	$b = 10$	A1ft	ft $18 - \text{their } 4 \times 2$ if B0M3
	<b>Alternative method 2</b>		
	$-\frac{1}{4}$ or $-0.25$	B1	gradient of $x + 4y = 74$ Do not allow embedded May be implied
	(gradient =) $\frac{-1}{\text{their } -\frac{1}{4}}$ or 4	M1	ft their $-\frac{1}{4}$ Only ft a non-zero numerical value Implied by $y = 4x + b$ or $a = 4$ (B1M1)
	Correct method for elimination of $y$ from $x + 4y = 74$ and $y = \text{their } 4x + b$	M1dep	eg $x + 4(4x + b) = 74$ or $17x + 4b = 74$
	Substitutes $x = 2$ into their equation	M1dep	eg $34 + 4b = 74$
	$b = 10$	A1ft	ft $18 - \text{their } 4 \times 2$ if B0M3

**Alternative method 3 and Additional Guidance continue on the next page**

<b>14 cont</b>	<b>Alternative method 3</b>		
	$-\frac{1}{4}$ or $-0.25$	B1	gradient of $x + 4y = 74$ Do not allow embedded May be implied
	(gradient =) $\frac{-1}{\text{their } -\frac{1}{4}}$ or 4	M1	ft their $-\frac{1}{4}$ Only ft a non-zero numerical value Implied by $y = 4x + b$ or $a = 4$ (B1M1)
	$(y =) \frac{74-2}{4}$ or $\frac{72}{4}$ or 18	M1	oe May be seen on diagram
	Correct method for elimination of $x$ from $x + 4y = 74$ and $y = \text{their } 4x + b$ <b>and</b> substitutes $y = \text{their } 18$	M1dep	eg $y = 4(74 - 4y) + b$ or $17y = 296 + b$ <b>and</b> $306 = 296 + b$ dep on M2
	$b = 10$	A1ft	ft 18 – their $4 \times 2$ if B0M3
	<b>Additional Guidance</b>		
$y = 4x + 10$ will gain full marks unless contradicted			
If an error is made in the constant term when rearranging $x + 4y = 74$ the B1 can still be awarded for gradient = $-\frac{1}{4}$ eg $y = -\frac{1}{4}x + 19$ and gradient = $-\frac{1}{4}$ is B1 (all other marks are possible)			
In alt 1 and alt 3 the mark for $y = 18$ will sometimes be the only mark awarded			

<b>15</b>	<b>Alternative method 1</b>			
	$wy = 8x - y$	$w = \frac{8x}{y} - 1$ or $w + 1 = \frac{8x}{y}$	M1	
	$wy + y = 8x$ or $y(w + 1) = 8x$	$\frac{w+1}{8x} = \frac{1}{y}$	M1dep	oe $y$ term(s) collected eg $-wy - y = -8x$ M2 $\frac{8x}{w+1}$ or $\frac{-8x}{-w-1}$ or $\frac{-8x}{-(w+1)}$
	$y = \frac{8x}{w+1}$ or $y = \frac{-8x}{-w-1}$ or $y = \frac{-8x}{-(w+1)}$		A1	oe eg $y = \frac{4x}{0.5w+0.5}$ Must have $y =$
	<b>Alternative method 2</b>			
	$y = \frac{8x}{w} - \frac{y}{w}$		M1	
	$y + \frac{y}{w} = \frac{8x}{w}$ or $y(1 + \frac{1}{w}) = \frac{8x}{w}$		M1dep	oe $y$ term(s) collected M2 $\frac{\frac{8x}{w}}{1 + \frac{1}{w}}$
	$y = \frac{\frac{8x}{w}}{1 + \frac{1}{w}}$		A1	oe Must have $y =$
	<b>Additional Guidance</b>			
	$y = \frac{8x}{w+1}$ in working with $\frac{8x}{w+1}$ on answer line etc			M2A1
Allow multiplications signs and 1s throughout				
$w = \frac{8x}{y} - \frac{y}{y}$ with no further simplification			M0	
Correct answer followed by incorrect further work			M2A0	

16(a)	$3^{-2b}$	B1	
	<b>Additional Guidance</b>		
16(b)	$5^{x+2}$	B1	
	<b>Additional Guidance</b>		
16(c)	$2^{3m}$	B1	
	<b>Additional Guidance</b>		
17(a)	$3x^2$ or $(-12x)$	M1	Attempt at $\frac{dy}{dx}$
	their $(3x^2 - 12x) = 0$	M1dep	Must have at least 2 terms for their $\frac{dy}{dx}$ The $= 0$ can be implied by sight of a correct non-zero solution to their $(3x^2 - 12x) = 0$
	$x = 4$ (and $x = 0$ )	A1ft	ft M2 if their $\frac{dy}{dx}$ is a 2-term quadratic
	$(4, -25)$ with correct expression for $\frac{dy}{dx}$ seen	A1	
	<b>Additional Guidance</b>		
	Condone $y = 3x^2 - 12x$ etc		M1
	Ignore working for second derivative or testing for minimum point		
	Stating $\frac{dy}{dx} = 0$ is not sufficient for second M mark but may be implied by correct solution(s) seen		

<b>17(b)</b>	<b>Alternative method 1</b>		
	$(-1)^3 - 6(-1)^2 + 7 = 0$ with no incorrect evaluations seen or $-1 - 6 + 7 = 0$	B1	Must have = 0
	<b>Alternative method 2</b>		
	$(x + 1)(x^2 - 7x + 7) = 0$ <b>and</b> $(x + 1) = 0$ <b>and</b> $x = -1$	B1	
	<b>Additional Guidance</b>		
	$(-1)^3 - 6(-1)^2 + 7$ or $-1 - 6 + 7$		B0
	Allow $-1^3$ or $(-1^3)$ for $(-1)^3$		
	Allow recovery of brackets for $(-1)^2$ eg1 $-1^3 - 6 \times -1^2 + 7 = 0$ eg2 $-1^3 - 6 \times -1^2 + 7 = -1 - 6 + 7 = 0$		B0 B1

<b>17(c)</b>	<b>Alternative method 1</b>		
	$(x - -1)$ or $(x + 1)$ seen	M1	
	$(x + 1)(x^2 - 7x + c)$	M1dep	$c$ can be any non-zero value Implied by $(x + 1)(x^2 + bx + c)$ <b>and</b> $b + 1 = -6$ or $b = -7$
	$x^2 - 7x + 7 (= 0)$	A1	
	$\frac{- -7 \pm \sqrt{(-7)^2 - 4 \times 1 \times 7}}{2 \times 1}$ or $\frac{7 \pm \sqrt{21}}{2}$	M1	oe eg $\frac{7}{2} \pm \sqrt{\frac{21}{4}}$  Correct attempt to solve their 3-term quadratic Allow recovery of brackets Allow $7^2$ for $(-7)^2$  Implied by correct solutions to their 3-term quadratic seen
	5.79 and 1.21 with $x^2 - 7x + 7 (= 0)$ seen	A1	Must both be to 2 dp
	<b>Alternative method 2</b>		
	$(x - -1)$ or $(x + 1)$ seen	M1	
	$x^2 - 7x \dots$ $x + 1 \overline{)x^3 - 6x^2 (+0x) + 7}$	M1dep	
	$x^2 - 7x + 7 (= 0)$	A1	
	$\frac{- -7 \pm \sqrt{(-7)^2 - 4 \times 1 \times 7}}{2 \times 1}$ or $\frac{7 \pm \sqrt{21}}{2}$	M1	oe eg $\frac{7}{2} \pm \sqrt{\frac{21}{4}}$  Correct attempt to solve their 3-term quadratic Allow recovery of brackets Allow $7^2$ for $(-7)^2$  Implied by correct solutions to their 3-term quadratic seen
	5.79 and 1.21 with $x^2 - 7x + 7 (= 0)$ seen	A1	Must both be to 2 dp

**Additional Guidance is on the next page**



		<b>Additional Guidance</b>
<b>17(c) cont</b>	Final A1 mark can be awarded if both answers seen in working with $x^2 - 7x + 7 (= 0)$ seen but only one answer is written on answer line	
	$(x + 1)$ followed by 5.79 and 1.21 without $x^2 - 7x + 7 (= 0)$ seen	M1MOA0 MOA0
	$(x - 1)$ instead of $(x + 1)$ can score a maximum of M0M0A0M1A0	
	T & I on the cubic equation	Zero

<b>18</b>	$x^2 + (2x)^2 = (4y)^2$ or $x^2 + 4x^2 = 16y^2$	M1	oe eg $5x^2 = (4y)^2$ Missing brackets may be recovered
	$x^2 = \frac{16}{5}y^2$ or $5x^2 = 16y^2$ or $x = \sqrt{\frac{16y^2}{5}}$ or $x\sqrt{5} = 4y$	M1	oe equation of the form $ax^2 = by^2$ or $cx = dy$ or $x = \sqrt{ky^2}$ eg $x^2 = 16y^2 \div 5$ ft if Pythagoras used with only error being missing brackets
	$2 \times \text{their } \frac{16}{5}y^2$ or their $16y^2 \div \frac{\text{their } 5}{2}$	M1dep	oe $2 \times \text{their } x^2$ or $2 \times (\text{their } x)^2$ dep on at least one M
	$\frac{32}{5}y^2$ or $6\frac{2}{5}y^2$ or $6.4y^2$	A1	
	<b>Additional Guidance</b>		
	$5x^2 = (4y)^2$ with no further work		M1M0M0
	$x^2 + 4x^2 = 4y^2$ with answer $\frac{8}{5}y^2$		M0M1M1A0
	$x^2 + 2x^2 = 16y^2$ with answer $\frac{32}{3}y^2$		M0M1M1A0
	$x^2 + 2x^2 = 4y^2$ with answer $\frac{8}{3}y^2$		M0M1M1A0
	$(2x)^2 = (4y)^2 - x^2$ $x^2 = 16y^2$ and $2 \times 16y^2 = 32y^2$		M1 M0M1A0
$\frac{32}{5}y^2$ followed by further work		M3A0	

<b>19(a)</b>	$k$	B1	
	<b>Additional Guidance</b>		
	$k = 0$ or $k = 1$ etc		B0

<b>19(b)</b>	$-k$	B1	
	<b>Additional Guidance</b>		
	$-k = 0$ or $-k = 1$ etc		B0

<b>19(c)</b>	$k^2 + \cos^2 \alpha = 1$ or $1 - k^2$	M1	oe eg $(1 + k)(1 - k)$
	$\sqrt{1 - k^2}$ or $\sqrt{(1 + k)(1 - k)}$	A1	
	<b>Additional Guidance</b>		
	Answer $-\sqrt{1 - k^2}$ or $\pm\sqrt{1 - k^2}$		M1A0
	Correct answer followed by incorrect further work		M1A0
	Answer $1 - k^2$		M1A0
	Allow $\cos^2 x$ or $\cos^2 \theta$ etc or $\cos^2$ or $c^2$ or $(\cos \alpha)^2$ for $\cos^2 \alpha$		
	Condone $\cos \alpha^2$ for $\cos^2 \alpha$		
$\cos(\sin^{-1} k)$		M0A0	

<b>20(a)</b>	Angle in a semicircle (is a right angle) or Angle at centre is $180^\circ$ , angle at circumference is half the angle at the centre ( $= 90^\circ$ ) or Angle at centre is $180^\circ$ , angle at centre is twice the angle at the circumference or Angle subtended at circumference by a diameter	<b>B1</b>	
	<b>Additional Guidance</b>		
	Do not allow half a circle to mean a semicircle		
	Allow extra words if not contradictory		
	eg1 Angle at the circumference in a semicircle		B1
	eg2 Angle inscribed in a semicircle		B1
	Angle subtended by a diameter (no mention of at circumference)		B0
	Angle in a hemisphere is 90		B0
	Angle at centre is 180		B0
	Angle at circumference is half the angle at the centre		B0
	2 chords on diameter meet at 90		B0
	Triangle in a semicircle always has a right angle		B0
	Angle in a semicircle is 180		B0
	Angle on a diameter is a right angle		B0
Because $AB$ is a diameter		B0	

<b>20(b)</b>	angle $ABE$ $= 90 - x$ <b>or</b> angle $CBE$ $= 90 + x$	angle $DEB = 90$ <b>and</b> angle $DCB = 90$	B1	
	angle $CDE = 90 - x$		B1dep	
	angle $CED = 90 - x$		B1dep	
	angle $DCE = 2x$ <b>and</b> all reasons given for their proof		B1dep	See guidance for acceptable wording for reasons
	<b>Additional Guidance</b>			
	To award a particular mark, all previous marks must have been awarded			
	First three B marks can be awarded with no or incorrect reasons			
	Do not mark any working on the diagram – statements are needed			
	Incorrect angles score B0			
	eg1 angle $ABE = 90 - x$ angle $DEC = 90 + x$			B1B0B0B0
	eg2 angle $ABE = 90 - x$ angle $CDE = 90 - x$ angle $DCE = 90 + x$			B1B1B0B0
	Angle $CDE$ and angle $CDA$ are the same angle etc			
	Angle $EBA$ and angle $ABE$ are the same angle etc			
	Condone $ABE$ for angle $ABE$ etc			
Do not allow angle $C$ for angle $DCE$ etc				
$CE$ must be proven to be a tangent if used in a response				
Reasons angle sum of triangle (is $180^\circ$ ) or angles in a triangle (add to $180^\circ$ ) or $180^\circ$ in a triangle (adjacent) angles on a (straight) line (add to $180^\circ$ ) or $180^\circ$ on a (straight) line exterior angle of triangle (= sum of opposite interior angles) (equal angles in an) isosceles (triangle) or $CD = CE$ (opposite angles in a) cyclic quadrilateral (add to $180^\circ$ ) exterior angle of cyclic quadrilateral (= opposite interior angle)			Degrees symbol may be omitted  Abbreviations are allowed eg quad for quadrilateral	

<b>21(a)</b>	(0, 8)	B1	
	<b>Additional Guidance</b>		
	Answer line takes precedence over working lines and diagram		
	Answer line blank with C labelled (0, 8) on diagram		B1
	Answer line blank with 8 written next to C on diagram		B0
	(8, 0)		B0
	Answer 8		B0

<b>21(b)</b>	$-x^2 - 2x + 4x + 8$	M1	Allow one error but no omission Must have an $x^2$ term Terms may be seen in a grid Implied by $-x^2 + 2x + k \quad k \neq 0$ or $ax^2 + 2x + 8 \quad a \neq 0$
	$-x^2 - 2x + 4x + 8$ or $-x^2 + 2x + 8$	A1	$-x^2 - 2x + 4x + 8$ but an error in any collection of terms is M1A0
	$-2x - 2 + 4$ or $-2x + 2$ or $-2(x - 1)$ or $2(1 - x)$	A1ft	oe ft their quadratic in $x$ with M1 awarded
	<b>Additional Guidance</b>		
	2 – 2x with final answer 2 (from substituting in $x = 0$ )		M1A1A0
	Condone $y = 2 - 2x$ or $f(x) = 2 - 2x$ in working for M1A1 If $(\frac{dy}{dx} \text{ or } f'(x) =) 2 - 2x$ on answer line also award final A1		
	$y = 2 - 2x$ or $f(x) = 2 - 2x$ on answer line		M1A1A0
	When marking (b), a maximum of M1A1A0 can be awarded from an expansion seen on the previous page if not contradicted by an expansion in (b) The final A1 must be seen in (b) eg1 (b) no expansion seen with an answer of $2x + 2$ At top of previous page $-x^2 + 2x + 8$ eg2 (b) no expansion seen with an answer of $-2x + 6$ In (a) $-x^2 + 2x + 4x + 8 = -x^2 + 6x + 8$		M1A1A0  M1A0A1ft
	Correct use of product rule and gradient function = $-2x + 2$		3 marks

<b>21(c)</b>	<b>Alternative method 1</b>		
	(gradient of curve at C =) 2	B1ft	Correct or ft their (b) when $x = 0$ May be implied
	$-\frac{1}{\text{their } 2}$ or $-\frac{1}{2}$	M1	oe ft their 2 Only ft a non-zero numerical value (gradient of normal =) $-\frac{1}{2}$ is B1M1
	$y = (\text{their } -\frac{1}{2})x + \text{their } 8$ or $y - \text{their } 8 = \text{their } -\frac{1}{2}(x - 0)$	M1dep	Must have used gradient of normal not gradient of tangent Correct or ft their 8 from (a) in the form (0, k)
	$0 = (\text{their } -\frac{1}{2})x + \text{their } 8$ or $0 - \text{their } 8 = \text{their } -\frac{1}{2}(x - 0)$ or $x = 16$	M1dep	Correct or ft their 8 from (a) in the form (0, k)
	$x = 16$ and $BD = 12$ and $AB = 6$ with correct method seen	A1	oe
	<b>Alternative method 2</b>		
	(gradient of curve at C =) 2	B1ft	Correct or ft their (b) when $x = 0$ May be implied
	$-\frac{1}{\text{their } 2}$ or $-\frac{1}{2}$	M1	oe ft their 2 Only ft a non-zero numerical value (gradient of normal =) $-\frac{1}{2}$ is B1M1
	$\frac{0 - \text{their } 8}{x - 0} = \text{their } -\frac{1}{2}$ or $x = -\text{their } 8 \div \text{their } -\frac{1}{2}$ or $x = 16$	M2dep	oe Correct or ft their 8 from (a) in the form (0, k)
	$x = 16$ and $BD = 12$ and $AB = 6$ with correct method seen	A1	oe
	<b>Additional Guidance</b>		

<b>22</b>	<b>Alternative method 1</b>		
	$(x - 2)^2 + (2x + 1 - 1)^2 = 16$	M1	oe Eliminates y
	$x^2 - 2x - 2x + 4 + 4x^2 = 16$ or $5x^2 - 4x - 12 (= 0)$	M1dep	oe Expands both brackets correctly
	$(5x + 6)(x - 2) (= 0)$ or $\frac{-4 \pm \sqrt{(-4)^2 - 4 \times 5 \times -12}}{2 \times 5}$	M1	oe eg $\frac{2}{5} \pm \sqrt{\frac{64}{25}}$ Correct attempt to solve their 3-term quadratic Allow recovery of brackets in formula Allow $4^2$ for $(-4)^2$ Implied by correct solutions to their 3-term quadratic seen
	$(x =) -1.2$ and $(x =) 2$ or $(x =) -1.2$ and $(y =) -1.4$ or $(x =) 2$ and $(y =) 5$ with $5x^2 - 4x - 12 (= 0)$ seen	A1	oe eg $(x =) -\frac{6}{5}$ and $(x =) 2$ with $5x^2 - 4x - 12 (= 0)$ seen
	$(-1.2, -1.4)$ and $(2, 5)$ with $5x^2 - 4x - 12 (= 0)$ seen	A1	oe eg $(-\frac{6}{5}, -\frac{7}{5})$ and $(2, 5)$ with $5x^2 - 4x - 12 (= 0)$ seen

<b>22</b>	<b>Alternative method 2</b>		
	$x^2 - 2x - 2x + 4 + y^2 - y - y + 1 = 16$	M1	oe Expands both brackets correctly
	$x^2 - 2x - 2x + 4 + (2x + 1)^2$ $- (2x + 1) - (2x + 1) + 1 = 16$ or $5x^2 - 4x - 12 (= 0)$	M1dep	oe Eliminates y
	$(5x + 6)(x - 2) (= 0)$ or $\frac{-4 \pm \sqrt{(-4)^2 - 4 \times 5 \times -12}}{2 \times 5}$	M1	oe eg $\frac{2}{5} \pm \sqrt{\frac{64}{25}}$ Correct attempt to solve their 3-term quadratic Allow recovery of brackets in formula Allow $4^2$ for $(-4)^2$ Implied by correct solutions to their 3-term quadratic seen
	$(x =) -1.2$ and $(x =) 2$ or $(x =) -1.2$ and $(y =) -1.4$ or $(x =) 2$ and $(y =) 5$ with $5x^2 - 4x - 12 (= 0)$ seen	A1	oe eg $(x =) -\frac{6}{5}$ and $(x =) 2$ with $5x^2 - 4x - 12 (= 0)$ seen
	$(-1.2, -1.4)$ and $(2, 5)$ with $5x^2 - 4x - 12 (= 0)$ seen	A1	oe eg $(-\frac{6}{5}, -\frac{7}{5})$ and $(2, 5)$ with $5x^2 - 4x - 12 (= 0)$ seen

**Alternative methods 3 and 4 and Additional Guidance continue on the next two pages**



<b>22</b>	<b>Alternative method 3</b>		
	$\left(\frac{y-1}{2}\right) - 2)^2 + (y-1)^2 = 16$	M1	oe Eliminates $x$
	$\left(\frac{y-1}{2}\right)^2 - 2\left(\frac{y-1}{2}\right) - 2\left(\frac{y-1}{2}\right) + 4$ $+ y^2 - y - y + 1 = 16$ or $5y^2 - 18y - 35 (= 0)$	M1dep	oe Expands $\left(\frac{y-1}{2}\right) - 2)^2$ and $(y-1)^2$ correctly
	$(5y+7)(y-5) (= 0)$ or $\frac{-18 \pm \sqrt{(-18)^2 - 4 \times 5 \times -35}}{2 \times 5}$	M1	oe eg $\frac{9}{5} \pm \sqrt{\frac{256}{25}}$ Correct attempt to solve their 3-term quadratic Allow recovery of brackets in formula Allow $18^2$ for $(-18)^2$ Implied by correct solutions to their 3-term quadratic seen
	$(y =) -1.4$ and $(y =) 5$ or $(x =) -1.2$ and $(y =) -1.4$ or $(x =) 2$ and $(y =) 5$ with $5y^2 - 18y - 35 (= 0)$ seen	A1	oe eg $(y =) -\frac{7}{5}$ and $(y =) 5$ with $5y^2 - 18y - 35 (= 0)$ seen
	$(-1.2, -1.4)$ and $(2, 5)$ with $5y^2 - 18y - 35 (= 0)$ seen	A1	oe eg $(-\frac{6}{5}, -\frac{7}{5})$ and $(2, 5)$ with $5y^2 - 18y - 35 (= 0)$ seen

**Alternative method 4 and Additional Guidance continue on the next page**

<b>22</b>	<b>Alternative method 4</b>		
	$x^2 - 2x - 2x + 4 + y^2 - y - y + 1 = 16$	M1	oe Expands both brackets correctly
	$\left(\frac{y-1}{2}\right)^2 - 2\left(\frac{y-1}{2}\right) - 2\left(\frac{y-1}{2}\right) +$ $4 + y^2 - y - y + 1 = 16$ or $5y^2 - 18y - 35 (= 0)$	M1dep	oe Eliminates $x$
	$(5y + 7)(y - 5) (= 0)$ or $\frac{-(-18) \pm \sqrt{(-18)^2 - 4 \times 5 \times -35}}{2 \times 5}$	M1	oe eg $\frac{9}{5} \pm \sqrt{\frac{256}{25}}$ Correct attempt to solve their 3-term quadratic Allow recovery of brackets in formula Allow $18^2$ for $(-18)^2$ Implied by correct solutions to their 3-term quadratic seen
	$(y =) -1.4$ and $(y =) 5$ or $(x =) -1.2$ and $(y =) -1.4$ or $(x =) 2$ and $(y =) 5$ with $5y^2 - 18y - 35 (= 0)$ seen	A1	oe eg $(y =) -\frac{7}{5}$ and $(y =) 5$ with $5y^2 - 18y - 35 (= 0)$ seen
	$(-1.2, -1.4)$ and $(2, 5)$ with $5y^2 - 18y - 35 (= 0)$ seen	A1	oe eg $(-\frac{6}{5}, -\frac{7}{5})$ and $(2, 5)$ with $5y^2 - 18y - 35 (= 0)$ seen
	<b>Additional Guidance</b>		
Answers only (no valid working)		Zero	
Both solutions from scale drawing		5 marks	
$(2, 5)$ is often seen without seeing any correct method		Zero	
Allow one miscopy for up to M3A0A0			

<b>23</b>	<b>Alternative method 1</b>		
	Replaces $\tan x$ with $\frac{\sin x}{\cos x}$ at least once in given expression	M1	eg $\frac{1}{\frac{\sin^2 x}{\cos^2 x}} - \frac{1}{\sin^2 x}$
	Correct steps leading to the single fraction $\frac{\cos^2 x - 1}{\sin^2 x}$ or $\frac{\cos^2 x - 1}{1 - \cos^2 x}$ or $\frac{1 - \sin^2 x - 1}{\sin^2 x}$ or $\frac{\cos^2 x - \cos^2 x - \sin^2 x}{\sin^2 x}$ or $\frac{-\sin^2 x}{\sin^2 x}$	M1dep	
	$\frac{\cos^2 x - 1}{\sin^2 x} = \frac{-\sin^2 x}{\sin^2 x} = -1$ or $\frac{\cos^2 x - 1}{1 - \cos^2 x} = -1$ or $\frac{1 - \sin^2 x - 1}{\sin^2 x} = -1$ or $\frac{-\sin^2 x}{\sin^2 x} = -1$	A1	Must see all steps leading to $-1$

**Alternative method 2 and Additional Guidance continue on the next page**

<b>23 cont</b>	<b>Alternative method 2</b>		
	Replaces $\tan x$ with $\frac{\sin x}{\cos x}$ at least once in given expression	M1	eg $\frac{\sin^2 x - \frac{\sin^2 x}{\cos^2 x}}{\sin^2 x \frac{\sin^2 x}{\cos^2 x}}$
	Correct steps leading to the single fraction $\frac{\sin^2 x (\cos^2 x - 1)}{\sin^4 x}$ or $\frac{-\sin^4 x}{\sin^4 x}$	M1dep	
	$\frac{\sin^2 x (\cos^2 x - 1)}{\sin^4 x} = \frac{-\sin^4 x}{\sin^4 x} = -1$ or $\frac{-\sin^4 x}{\sin^4 x} = -1$	A1	Must see all steps leading to $-1$
	<b>Additional Guidance</b>		
	Allow $\cos^2 \theta$ etc or $\cos^2$ or $c^2$ or $(\cos x)^2$ for $\cos^2 x$ etc		
	Condone $\cos x^2$ for $\cos^2 x$ etc		
	Only substituting values for $x$		Zero
	$\frac{\cos^2 x - 1}{\sin^2 x}$ etc with no working		Zero
	Alt 2 $\frac{\sin^2 x \cos^2 x - \sin^2 x}{\sin^4 x}$ with no further working		M1M0A0
Any fully correct response that shows how the given expression is equal to $-1$ is awarded 3 marks  eg $\frac{1}{\frac{\sin^2 x}{\cos^2 x}} - \frac{1}{\sin^2 x} = \frac{\cos^2 x}{\sin^2 x} - \frac{1}{\sin^2 x} = \frac{1 - \sin^2 x}{\sin^2 x} - \frac{1}{\sin^2 x}$  $= \frac{1}{\sin^2 x} - \frac{\sin^2 x}{\sin^2 x} - \frac{1}{\sin^2 x} = -1$		3 marks	
$\cot^2 x - \operatorname{cosec}^2 x = -1$		3 marks	

<b>24</b>	<b>Alternative method 1</b>		
	$12(x^2 - 5x) \dots$ or $12(x - 2.5)^2 \dots$	M1	oe eg $12\{(x^2 - 5x) \dots\}$ or $12(x^2 - 5x \dots)$
	$12\{(x - 2.5)^2 - 2.5^2\} \dots$ or $12(x - 2.5)^2 - 75 \dots$	M1dep	oe eg $12\{(x - 2.5)^2 - 2.5^2 \dots\}$
	$12(x - 2.5)^2 - 12 \times 2.5^2 + 5$ or $12(x - 2.5)^2 - 70$	M1dep	oe eg $12(x - 2.5)^2 - 12 \times 2.5^2 + 12 \times \frac{5}{12}$
	$12\left(\frac{2x-5}{2}\right)^2 - 12 \times 2.5^2 + 5$	M1dep	oe eg $12\left(\frac{2x-5}{2}\right)^2 - 12 \times 2.5^2 + 12 \times \frac{5}{12}$
$3(2x - 5)^2 - 70$ or $a = 3 \quad b = 2 \quad c = -5 \quad d = -70$ or $3(5 - 2x)^2 - 70$ or $a = 3 \quad b = -2 \quad c = 5 \quad d = -70$	A1	oe	

<b>Alternative method 2</b>		
$3(4x^2 - 20x) \dots$ or $3(2x - 5)^2 \dots$	M1	oe eg $3\{(4x^2 - 20x) \dots\}$ or $3(4x^2 - 20x \dots)$
$3\{(2x - 5)^2 - 5^2\} \dots$ or $3(2x - 5)^2 - 75 \dots$	M1dep	oe eg $3\{(2x - 5)^2 - 5^2 \dots\}$
$3\{(2x - 5)^2 - 5^2\} + 5$	M1dep	oe eg $3\{(2x - 5)^2 - 5^2 + \frac{5}{3}\}$
$3(2x - 5)^2 - 3 \times 5^2 + 5$	M1dep	oe eg $3(2x - 5)^2 - 3 \times 5^2 + 3 \times \frac{5}{3}$
<b>24</b> $3(2x - 5)^2 - 70$ or $a = 3 \quad b = 2 \quad c = -5 \quad d = -70$ or $3(5 - 2x)^2 - 70$ or $a = 3 \quad b = -2 \quad c = 5 \quad d = -70$	A1	oe
<b>Additional Guidance</b>		
For M marks 2.5 may be seen as $\frac{5}{2}$		
For M marks $(x - 2.5)^2$ may be replaced by $(2.5 - x)^2$ etc		
Expansion of given form followed by trial and improvement eg1 $3(2x - 5)^2 - 70$ (or $a = 3 \quad b = 2 \quad c = -5 \quad d = -70$ ) eg2 Not fully correct		5 marks Zero