

A Level Mathematics B (MEI) H640/03 Pure Mathematics and Comprehension Sample Question Paper

Version 2.1

Date – Morning/Afternoon

Time allowed: 2 hours

You must have:

- Printed Answer Booklet
- the Insert

You may use:

- a scientific or graphical calculator



INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total number of marks for this paper is **75**.
- The marks for each question are shown in brackets [].
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is used. You should communicate your method with correct reasoning.
- The Printed Answer Booklet consists of **20** pages. The Question Paper consists of **12** pages.

Formulae A Level Mathematics B (MEI) (H640)

Arithmetic series

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a+(n-1)d\}$$

Geometric series

$$S_n = \frac{a(1-r^n)}{1-r}$$

$$S_\infty = \frac{a}{1-r} \quad \text{for } |r| < 1$$

Binomial series

$$(a+b)^n = a^n + {}^n C_1 a^{n-1}b + {}^n C_2 a^{n-2}b^2 + \dots + {}^n C_r a^{n-r}b^r + \dots + b^n \quad (n \in \mathbb{N}),$$

$$\text{where } {}^n C_r = {}_n C_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

Differentiation

f(x)	f'(x)
tan kx	k sec ² kx
sec x	sec x tan x
cot x	-cosec ² x
cosec x	-cosec x cot x

$$\text{Quotient Rule } y = \frac{u}{v}, \quad \frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Integration

$$\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + c$$

$$\int f'(x)(f(x))^n dx = \frac{1}{n+1}(f(x))^{n+1} + c$$

$$\text{Integration by parts } \int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

Small Angle Approximations

$$\sin \theta \approx \theta, \quad \cos \theta \approx 1 - \frac{1}{2}\theta^2, \quad \tan \theta \approx \theta \quad \text{where } \theta \text{ is measured in radians}$$

Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad (A \pm B \neq (k + \frac{1}{2})\pi)$$

Numerical methods

Trapezium rule: $\int_a^b y \, dx \approx \frac{1}{2} h \{ (y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1}) \}$, where $h = \frac{b-a}{n}$

The Newton-Raphson iteration for solving $f(x) = 0$: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A)P(B | A) = P(B)P(A | B) \quad \text{or} \quad P(A | B) = \frac{P(A \cap B)}{P(B)}$$

Sample Variance

$$s^2 = \frac{1}{n-1} S_{xx} \quad \text{where} \quad S_{xx} = \sum (x_i - \bar{x})^2 = \sum x_i^2 - \frac{(\sum x_i)^2}{n} = \sum x_i^2 - n\bar{x}^2$$

Standard deviation, $s = \sqrt{\text{variance}}$

The Binomial Distribution

If $X \sim B(n, p)$ then $P(X = r) = {}^n C_r p^r q^{n-r}$ where $q = 1 - p$


Mean of X is np

Hypothesis testing for the mean of a Normal distribution

If $X \sim N(\mu, \sigma^2)$ then $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ and $\frac{\bar{X} - \mu}{\sigma / \sqrt{n}} \sim N(0, 1)$

Percentage points of the normal distribution

p	10	5	2	1
z	1.645	1.960	2.326	2.576

**Kinematics**

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2} at^2$$

$$s = \frac{1}{2}(u + v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2} at^2$$

Motion in two dimensions

$$\mathbf{v} = \mathbf{u} + \mathbf{a}t$$

$$\mathbf{s} = \mathbf{u}t + \frac{1}{2} \mathbf{a}t^2$$

$$\mathbf{s} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$$

$$\mathbf{s} = \mathbf{v}t - \frac{1}{2} \mathbf{a}t^2$$

Answer **all** the questions

Section A (60 marks)

1 Express $\frac{2}{x-1} + \frac{5}{2x+1}$ as a single fraction. [2]

2 Find the first four terms of the binomial expansion of $(1-2x)^{\frac{1}{2}}$.

State the set of values of x for which the expansion is valid. [4]

3 Show that points A (1, 4, 9), B (0, 11, 17) and C (3, -10, -7) are collinear. [4]

4 Show that $\sum_{r=1}^4 \ln \frac{r}{r+1} = -\ln 5$. [3]

5 In this question you must show detailed reasoning.

Fig. 5 shows the circle with equation $(x-4)^2 + (y-1)^2 = 10$.

The points $(1, 0)$ and $(7, 0)$ lie on the circle. The point C is the centre of the circle.




Fig. 5

Find the area of the part of the circle below the x -axis.

[5]

- 6 Fig. 6 shows the curve with equation $y = x^4 - 6x^2 + 4x + 5$.




Fig. 6

Find the coordinates of the points of inflection.

[5]

- 7 By finding a counter example, disprove the following statement.

If p and q are non-zero real numbers with $p < q$, then $\frac{1}{p} > \frac{1}{q}$.

[2]

- 8 In Fig. 8, OAB is a thin bent rod, with $OA = 1$ m, $AB = 2$ m and angle $OAB = 120^\circ$. Angles θ , ϕ and h are as shown in Fig. 8.




Fig. 8

- (a) Show that $h = \sin \theta + 2 \sin(\theta + 60^\circ)$. [3]

The rod is free to rotate about the origin so that θ and ϕ vary. You may assume that the result for h in part (a) holds for all values of θ .

- (b) Find an angle θ for which $h = 0$. [5]

- 9 (a) Express $\cos \theta + 2 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $0 < \alpha < \frac{1}{2}\pi$ and R is positive and given in exact form. [4]

The function $f(\theta)$ is defined by $f(\theta) = \frac{1}{(k + \cos \theta + 2 \sin \theta)}$, $0 \leq \theta \leq 2\pi$, k is a constant.

- (b) The maximum value of $f(\theta)$ is $\frac{(3 + \sqrt{5})}{4}$.
Find the value of k . [3]

10 The function $f(x)$ is defined by $f(x) = x^4 + x^3 - 2x^2 - 4x - 2$.

- (a) Show that $x = -1$ is a root of $f(x) = 0$. [1]

- (b) Show that another root of $f(x) = 0$ lies between $x = 1$ and $x = 2$. [2]

- (c) Show that $f(x) = (x + 1)g(x)$, where $g(x) = x^3 + ax + b$ and a and b are integers to be determined. [3]

- (d) Without further calculation, explain why $g(x) = 0$ has a root between $x = 1$ and $x = 2$. [1]

- (e) Use the Newton-Raphson formula to show that an iteration formula for finding roots of $g(x) = 0$ may be written

$$x_{n+1} = \frac{2x_n^3 + 2}{3x_n^2 - 2}.$$

Determine the root of $g(x) = 0$ which lies between $x = 1$ and $x = 2$ correct to 4 significant figures. [3]

11 The curve $y = f(x)$ is defined by the function $f(x) = e^{-x} \sin x$ with domain $0 \leq x \leq 4\pi$.

(a) (i) Show that the x -coordinates of the stationary points of the curve $y = f(x)$, when arranged in increasing order, form an arithmetic sequence.

(ii) Show that the corresponding y -coordinates form a geometric sequence. **[9]**

(b) Would the result still hold with a larger domain? Give reasons for your answer. **[1]**

Answer **all** the questions

Section B (15 marks)

The questions in this section refer to the article on the Insert. You should read the article before attempting the questions.

- 12** Explain why the smaller regular hexagon in **Fig. C1** has perimeter 6. [1]
- 13** Show that the larger regular hexagon in **Fig. C1** has perimeter $4\sqrt{3}$. [3]
- 14** Show that the two values of b given on line 36 are equivalent. [3]

- 15 Fig. 15 shows a unit circle and the escribed regular polygon with 12 edges.




Fig. 15

- (a) Show that the perimeter of the polygon is $24 \tan 15^\circ$. [2]
- (b) Using the formula for $\tan(\theta - \phi)$ show that the perimeter of the polygon is $48 - 24\sqrt{3}$. [3]
- 16 On a unit circle, the inscribed regular polygon with 12 edges gives a lower bound for π , and the escribed regular polygon with 12 edges gives an upper bound for π .

Calculate the values of these bounds for π , giving your answers:

- (i) in surd form
- (ii) correct to 2 decimal places. [3]

END OF QUESTION PAPER

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 9EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.